

## SCROPHULARINEÆ.

*Limnophila serrata*, Gaud.

*Limnophila serrata*, Gaud. ; Benth., Fl. Austr., iv. p. 490 ; Miq., Fl. Ind. Bat., ii. p. 680 ; Nadeaud, Enum. Pl. Tahiti, p. 59.

*Limnophila fragrans*, Seem., Fl. Vit., p. 180.

*Ruellia fragrans*, Forst., Fl. Ins. Austr. Prodr., p. 44.

ADMIRALTY ISLANDS.—Fijis, Tahiti, and North Australia, and Gaudichaud's original was from the Marianne Islands. It grows in swamps and stagnant pools on the coast. *Limnophila* is a genus of herbaceous marsh or aquatic plants, and comprises about twenty species spread over the warmer parts of the Old World.

*Limnophila roxburghii*, G. Don.

*Limnophila roxburghii*, G. Don ; Hook. f., Fl. Brit. Ind., iv. p. 265.

*Limnophila menthastrum*, Benth. ; Miq., Fl. Ind. Bat., ii. p. 680.

*Adenosma triflora*, Seem., Fl. Vit., p. 284, non Nees.

ADMIRALTY ISLANDS.—North India, China, Formosa, and the Philippines, southward to Java, the Fijis and Samoa.

## ACANTHACEÆ.

*Lepidagathis hyalina*, Nees.

*Lepidagathis hyalina*, Nees. ; Miq., Fl. Ind. Bat., ii. p. 815.

ADMIRALTY ISLANDS.—India and China to the South-eastern Moluccas.

*Hemigraphis reptans*, T. And.

*Ruellia reptans*, Forst., Fl. Ins. Austr. Prodr., p. 44 ; Seem., Fl. Vit., p. 183.

ADMIRALTY ISLANDS.—Southern Moluccas ; New Hebrides.

## VERBENACEÆ.

*Callicarpa cumingiana*, Schauer ?

*Callicarpa cumingiana*, Schauer ? Miq., Fl. Ind. Bat., ii. p. 887.

ADMIRALTY ISLANDS.—The typical *Callicarpa cumingiana* is from the Philippine Islands.

*Clerodendron inerme*, R. Br.

*Clerodendron inerme*, R. Br. ; Seem., Fl. Vit., p. 188 ; A. Gr. in Proc. Amer. Acad., vi. p. 50.

*Volkameria inermis*, Linn., Fl. Zeyl., p. 231.

ADMIRALTY ISLANDS.—A common seaside shrub in Tropical Asia, Australia, and Western Polynesia.