

	<i>T. dimidiata</i> , Scacchi = <i>T. cawlidia</i> and <i>T. urna</i> <i>antiqua</i> , Risso.	ranean ; Straits of Samos, &c., Ægean Sea ; off Gomera, Teneriffe ; off Gue- taria, Atlantic Coast of Spain ; off Madeira ; Canary Islands ; Adventure Bank. <i>Fossil</i> : Upper Tertiaries, Nice ; South of France ; Italy ; Rhodes ; Malta ; Austria ; &c.
	Sub-genus <i>Cistella</i> , Gray.	
	Loop two-lobed, a single median sub-marginal septum.	
20 to 45	<i>Cistella cistellula</i> , S. Wood, Ann. and Mag. Nat. Hist., vol. vi. p. 5, 1840. (Mr Dall considers <i>T. lunifera</i> , Philippi to be a synonym?). This is a minute but well-characterised species.	Norway ; Shetland ; North-West shore of Europe ; Zet- land ; near Weymouth, S. coast of England ; off Port en Bessin, Normandy (E. Desl.) ; Guernsey ; Cape Breton, France ; Mediterranean. <i>Fossil</i> : Upper Tertiaries, Coralline Crag, England ; Sicily ; Austria ; Rhodes.
30 to 130	<i>Cistella neapolitana</i> , Scacchi, Osser. Zool., vol. ii. p. 18, 1833 = <i>Argiope forbesi</i> , Dav. A good and abundant species.	Mediterranean ; Adriatic ; Canary Islands. <i>Fossil</i> : Upper Tertiaries, South of Italy ; Rhodes ; Austria. Straits of Messina, Medi- terranean.
150	(?) <i>Cistella neapolitana</i> , var. <i>biplicata</i> , Sequenza, Reale Acad. delle Scienze Fisiche e Matematica, 1876. This shell, of which I have seen the type and only specimen, seems to me to be very closely related to <i>C. neapolitana</i> . Dr Gwyn Jeffreys, Proc. Zool. Soc., 1878, refers it to <i>A. cuneata</i> , Risso.	
70 to 250	<i>Cistella barrettiana</i> , Dav., Proc. Zool. Soc., p. 103, pl. xii. fig. 3, a, b, (?) = <i>Argiope antillarum</i> , Crosse and Fischer, Journ. de Conch., p. 270, pl. xiii. fig. 7, 1866. Mr Crosse, in a letter to me (24th March 1870), writes that my <i>Arg. barrettiana</i> and his <i>Arg. antillarum</i> may eventually turn out to be one species.	North-East coast of Jamaica (Barrett) ; Guadalupe ; off Rio de Janeiro, 70 fms. (Rathborn).
(?)	<i>Cistella barrettiana</i> , var. <i>lutea</i> , Dall, Bull. Mus. Comp. Zool., 1871.	Tortugas.
60 to	(?) <i>Cistella woodwardiana</i> , Dav., Proc. Zool. Soc., p. 103, pl. xii. fig. 4, a-c, 1866.	Jamaica (Barrett).
28 to 200	<i>Cistella cuneata</i> , Risso, Faun. Europ. Merid., p. 388, fig. clxxix., 1826 = <i>T. soldaniana</i> , Risso = <i>T. pera</i> , Kuster. A good species.	Mediterranean ; Adriatic ; Canary Islands. <i>Fossil</i> : Upper Tertiaries, South of Italy ; Austria ; Rhodes.
200 to 250	(?) <i>Cistella schrammi</i> , Crosse and Fischer, Journ. de Conch., vol. xiv. p. 269, pl. viii. fig. 6, 1866. This species requires more study.	Guadalupe, West Indies.