

	1	2	3	4	5	6	7
	North Atlantic.	South Atlantic.	South Indian Ocean.	Australasia.	South Pacific.	North Pacific.	Eastern Asia.
POLYCOPEIDÆ.							
<i>Polycope cingulata</i> , n. sp.,							
<i>farus</i> , n. sp.,				+			
<i>orbicularis</i> , G. O. Sars,	+	+	+	+			
CYTHERELLIDÆ.							
<i>Cytherella cavernosa</i> , G. S. Brady,				+			
<i>cingulata</i> , G. S. Brady,				+			+
<i>cribrosa</i> , n. sp.,				+			
<i>dromedaria</i> , n. sp.,		+					
<i>irregularis</i> , n. sp.,	+						
<i>lata</i> , n. sp.,	+	+		+			
<i>latimarginata</i> , n. sp.,				+			
<i>polita</i> , G. S. Brady,		+		+			
<i>pulchra</i> , G. S. Brady,	+	+		+			
<i>punctata</i> , G. S. Brady,		+		+	+		
<i>semitalis</i> , G. S. Brady,				+			
<i>truncata</i> , G. S. Brady,				+			
<i>venusta</i> , n. sp.,						+	

LIST OF DREDGINGS AND TOW-NET GATHERINGS EXAMINED, WITH
THE SPECIES OF OSTRACODA FOUND IN EACH.

Vigo Bay. 11 fathoms. Mud from anchor. May 21, 1876.

Paracypris polita, G. O. Sars.
Cythere stimpsoni, G. S. Brady.
 tenera, G. S. Brady.
Loxoconcha guttata (Norman).
 Cytheropteron intermedium, G. S. Brady.
Paradoxostoma ensiforme, G. S. Brady.
Polycope orbicularis, G. O. Sars.

Off Gomera, Canaries. 620 fathoms. Sandy mud and shells. February 12, 1873.

Macrocypris canariensis, n. sp.
Bairdia, sp.