

STATION 151.—Off Heard Island. 75 fathoms. Mud. February 7, 1874.

Bairdia simplex, n. sp.
Cythere kerguelenensis, n. sp.
wyville-thomsoni, n. sp.
foveolata, n. sp.
Xestoleberis setigera, n. sp.
Cytheropteron assimile, n. sp.
Cytherideis lævata, n. sp.
Sclerochilus contortus (Norman).

STATION 158.—Lat. 50° 1' S., long. 123° 4' E. Tow-net. March 7, 1874.

Halocypris atlantica, Lubbock.

STATION 159.—Lat. 47° 25' S., long. 130° 12' E. Tow-net. March 12, 1874.

Halocypris atlantica, Lubbock.
brevirostris, Dana.

STATION 160.—Lat. 42° 42' S., long. 134° 10' E. 2600 fathoms. Bottom temperature, 0°·2 C. Red clay. March 13, 1874.

Cytheropteron abyssorum, n. sp.

STATION 161.—Off entrance to Port Philip, South Australia. 38 fathoms. Sand. April 1, 1874.

Philomedes wyville-thomsoni, n. sp.

STATION 162.—Off East Moncœur Island, Bass Strait. 38 to 40 fathoms. Sand. April 2, 1874.

Pontocypris faba (?) (Reuss).
Macrocypris maculata, G. S. Brady.
Aglaia (?), *pusilla*, n. sp.
Bythocypris reniformis, n. gen. and sp.
Bairdia villosa, n. sp.
amygdaloides, G. S. Brady.
foveolata, G. S. Brady.
victrix, G. S. Brady.