

with irregular transverse brown markings, and in some with brown centres. Scapulars tawny rufous and pure rufous, irregularly marked with brown zig-zags, several with a subterminal black mark and a terminal tawny rufous spot. Wing-coverts chestnut, with brown vermiculations; the greater coverts with large, pure white, ovoid terminal drops, mostly situated on the outer webs, and bounded above and on the inner margins with black. Quills brown on the inner webs, pale rufous on the outer, with obscure pale brown marks. Tertiaries pale rufous on the outer webs, much indented with the brown of the inner webs. Innermost tertiaries pale rufous on both webs, transversely marked with cloudy brown. Tips of several of the inner webs of the quills pale rufous. Rectrices pale rufous, traversed by several bars of rufous brown, each bar being bounded above and below by a brown line. Penultimate pair of rectrices with rufous spots, rather than bands. Exterior short pair with almost pure white spots on the inner web. Chin and throat dingy rufous, intersected by brown zig-zag lines. A gular collar, formed of broad feathers with a broad subterminal pure white band, bounded below and above by a narrow dark brown band. Feathers of the upper breast dingy rufous freckled brown, pale tawny rufous at the base, with brown transverse lines and traces of white spots on the margins of some of the webs. Concealed lower breast and flank-plumes mostly pure white, with dark grey markings near the base, and terminated with rufous freckled with brown. Abdominal feathers pale albescent rufous, striated with brown, and some marked with pure white. Feathers of the ventral region isabelline white, with small obscure pale tawny rufous margins. Under tail-coverts isabelline, the longer with subterminal marks of brown and rusty. Under wing-coverts greyish white, banded with rufous brown. Thigh-coverts dark rufous brown. Stiff nasal plumes rufous, tipped brown. Wing 6 inches, tail 4.5, tarsus 0.68, middle toe 0.75, bill (from forehead) 1.12, width of gape 1.37. The tarsus is feathered anteriorly for about one-fourth of its length; the bill is massive.

14. *Eudynamis mindanensis*, Linn.

*Cuculus mindanensis*, Linn. Syst. Nat., vol. i. p. 169, No. 3, 1766.

*Eudynamis mindanensis*, Walden, Trans. Zool. Soc., vol. ix. p. 162; Proc. Zool. Soc., 1877, pp. 543, 691, 823; 1878, p. 946; 1879, p. 70.

[No. 392, ♂.]  
 No. 393, ♂. } Malanipa. Eyes red.]  
 No. 394, ♀.]

Malanipa is a new locality for this species; but, from its proximity to Mindanao, the examples may be considered typical.

The two males are in full black plumage, with a green gloss. They have, however, a shorter wing than Guimaras examples, the only other Philippine examples I have been