

26. *Hypsipetes philippensis*, Gm.

*Turdus philippensis*, Gm., S. N., vol. i. p. 814, No. 40 (1788).

*Hypsipetes philippensis*, Walden, Trans. Zool. Soc., vol. ix. p. 191; Proc. Zool. Soc., 1877, pp. 546, 696, 761, 827; 1878, pp. 285, 342, 380.

[No. 378, ♂. Zebu. Eyes hazel.]

27. *Hypsipetes rufigularis*, Sharpe.

*Hypsipetes rufigularis*, Sharpe, Trans. Linn. Soc., ser. 2, Zool., vol. i. p. 335; Walden, Proc. Zool. Soc., 1877, pp. 546, 827; 1878, p. 949; 1879, p. 72.

[No. 421, ♂.]

No. 422, ♂.] Pasananca. Eyes brown or hazel.]

No. 423, ♀.]

The female is somewhat smaller, with a shorter bill. An addition to the fauna of Mindanao.

28. *Orthotomus castaneiceps*, Walden.

*Orthotomus castaneiceps*, Walden, Ann. Nat. Hist., ser. 4, vol. i. p. 252, "Guimaras," 1st Oct. 1872; Walden, Trans. Zool. Soc., vol. ix. p. 195; Proc. Zool. Soc., 1878, p. 546; 1878, p. 286; Sharpe, Trans. Linn. Soc., ser. 2, Zool., vol. i. p. 337; id. Ibis, 1877, p. 114.

The type was a male; but this Panay female does not differ.

[No. 359, ♀. Iloilo. Eyes brown. Stomach had insects.]

29. *Calobates melanope*, Pallas.

*Motacilla melanope*, Pallas, Reisen Russischen Reichs, vol. iii. p. 696, No. 16, 1776.

*Calobates melanope*, Walden, Trans. Zool. Soc., vol. ix. p. 196; Proc. Zool. Soc., 1877, pp. 546, 697, 761; 1878, p. 343.

[No. 432, ♀. Pasananca.]

A female, with greyish brown upper plumage; tail, 3·56. Recorded for the first time from Mindanao.

30. *Corydalla lugubris*, Walden.

*Corydalla lugubris*, Walden, Trans. Zool. Soc., vol. ix. p. 198, 1875; Proc. Zool. Soc., 1877, pp. 547, 696, 762, 828; 1878, pp. 286, 380, 710.

[No. 357, ♂. Iloilo. Eyes hazel. Stomach had seeds.]

Panay is an additional locality.

This example differs from the type in having the tertaries and most of the wing-coverts broadly margined externally with pale rufous, not albescent. The pectoral feathers are tinged with pale rufous; and the brown central streaks are much broader than in the type. The flanks are washed with pale rufous.