

31. *Dicaeum mindanense*, Tweedd. (Pl. V. fig. 1).*Dicaeum mindanense*, Tweeddale, Proc. Zool. Soc., 1877, p. 547.

[No. 433, ♂. Pasanauca.]

♂. Above, cheeks and ear-coverts fuliginous olive-green. Below greyish white, most grey on the breast; quills dark brown, narrowly margined with olive-green on outer webs; tertiaries with the olive-green margins broader; rectrices dark brown, faintly edged with olive; axillaries and under wing-coverts white; feet and bill black. Wing, 1.0; tail, 1.1; bill, from forehead, 0.59; tarsus, 1.5. The bill is not normal, being longer than the head.

This species is a representative form of *D. hypoleucum*, ex Basilan, which is black above and larger.

32. *Nectarophila julia*, Tweedd. (Pl. V. figs. 2, 3).*Nectarophila julia*, Tweeddale, Proc. Zool. Soc., 1877, p. 547; 1878, p. 951.

[No. 402, ♂. }
 No. 403, ♂. } Malanipa. Eyes hazel. Were quite numerous on the island, generally
 No. 404, ♀. } about the tops of the high trees.]

♂. Forehead, entire head, and nape, brilliant metallic coppery green; chin, throat, and upper breast brilliant metallic amethystine; space before the eyes black; cheeks and ear-coverts brown; lesser wing-coverts, uropygium, and upper tail-coverts brilliant metallic emerald-green; breast-feathers deep yellow, with orange-coloured tips; abdomen, vent, and under tail-coverts yellowish olive-green; under wing-coverts pale yellow and white; axillaries pale yellow; back and major wing-coverts and scapulars maroon-red; quills dark brown, externally margined with yellowish ruddy-brown; rectrices black; bill and feet black. The second example has no orange edging to the pectoral feathers.

♀. Dull olive-green above, a narrow line passing from the base of bill over the eye pale yellow; chin, throat, abdomen, vent, and under tail-coverts pale yellow; breast the same, tinged with grey; minor wing-coverts olive-green, some tipped and centred with ruddy brown; quills dark brown, margined with ruddy brown; greater wing-coverts and scapulars ruddy brown, mixed with olive-green; inner edges of some of the quills margined with white; axillaries and under wing-coverts pale yellow and white; cheeks pale yellow; ear-coverts olive-green; rectrices black, middle pair narrowly, the others broadly tipped with white, outer pair for nearly half their length white; bill and feet black.

	Wing.	Tail.	Tarsus.	Bill, from gape.
♂	1.94	1.35	0.5	0.75
♀	1.87	1.25	0.5	0.62

A true *Nectarophila*. The figures (Pl. V.) are taken from specimens 403 and 404.