

Eyes red. Stomach contained fruit.]

As I have already stated (Ibis, 1871, p. 360), this bird has, in my opinion, no relationship to the Fringillidæ, but is merely an exaggerated form of the same type as *Hemignathus* and its allied genera.

5. *Drepanis coccinea*, Merr.

*Mellisuga coccinea*, Merram, Av. Descr. et Ic., p. 14, pl. iv., 1786.

*Drepanis coccinea*, Cassin, Zool. U. S. Expl. Exp., p. 177; Sci., Proc. Zool. Soc., p. 347.

[No. 521. Female adult. }  
 No. 522. Young female. } Hilo.  
 No. 523. " " }

Eyes red; bill and legs orange. Stomach contained insects and seeds.]

6. *Drepanis sanguinea*, Gm.

*Certhia sanguinea*, Gm., S. N., vol. i. p. 479.

*Drepanis sanguinea*, Bp., Consp., vol. i. p. 404; Sci., Proc. Zool. Soc., 1878, p. 347.

[No. 530. Female. Hilo.]

7. *Drepanis flava*, Bloxham.

*Drepanis flava*, Blox., Voy. Blonde, p. 249; Dole, Proc. Boston Soc., N. H., vol. xii. p. 298;  
 Sci., Proc. Zool. Soc., 1878, p. 348.

[No. 525. Male. }  
 No. 526. Male. } Hilo, Owhyhee.  
 No. 527. Female. }  
 No. 533. Female. }

Eyes black. Stomachs had seeds and insects.]

The females are much less yellowish than the males, especially below.

8. *Munia nitoria*, Temm..

*Munia nitoria*, Walden, Trans. Zool. Soc., vol. viii. p. 73; Sci., Proc. Zool. Soc., 1878,  
 p. 348.

[No. 543. Male. Sparrow, introduced; native who shot it said he had never seen one before.]

An imperfect skin of this species (or of the nearly allied *M. punctularia*), no doubt introduced.

Professor Baird a short time ago sent me a skin of a young or female of another Eastern Finch from the Sandwich Islands (probably *Munia malabarica*) for determination; also without doubt introduced.