No. 509. Female. North Pacific.

Eyes brown; bill black. Stomach empty. Shot on April 1, 1875, by Lord Campbell, with the Henry rifle, while on the wing. We were just north of the tropic; but this bird, as well as another Albatross, were seen some days before we had passed out of the tropics.

No. 511. Male. North Pacific.

Eyes brown; feet and bill dark or nearly black. Stomach had cuttlefish. Caught with a hook by Pearcey on April 7, 1875.

We first met this bird at sea in lat. 19° 25' N., in April 1875. It was after that a constant attendant on the ship, during the whole of our cruise in the North Pacific, till we reached 26° 11' N. of the Sandwich Islands.

It was occasionally in great numbers, and was frequently and easily taken with the hook at the stern of the ship.]

21. Diomedea melanophrys, Temm.

Diomedea melanophrys, Temm., PL Col., p. 456; Couce, Pr. Ac. Phil., 1866, p. 181; Sharpe, Zool Kerg., Birds, p. 46; Salv., Proc. Zool Soc., 1878, p. 740.

Male.

Shot at sea, July 8, 1874. Female.

Male.

Christmas Harbour, Kerguelen. Female.

This bird as well as Diomedea chlororhyncha is called the 'Molly Mauk' by the sailors and sealers. It was a frequent attendant on the ship, especially in the region of the southerly winds.

We did not notice it south of Heard Island. We never saw any nests of this bird.]

22. Diomedea fuliginosa, Gm.

Diomedea fuliginosa, Gm., Syst. Nat., vol. i. p. 595; Sharpe, Zool. Kerg., Birds, p. 48; Sulv., Proc. Zool. Soc., 1878, p. 740.

Phæbetria fuliginosa, Coues, Pr. Ac. Phil, 1866, p. 186; Coues and Kidder, Bull. Nat. Mus., No. 2, p. 21.

Male. Ice Barrier, February 16, 1874.

Female. Royal Sound, Kerguelen, January 17.

We met with this bird, the 'Piv' or 'Piew' of the scalers, all over the Southern and Antarctic Oceans, down to and even among the Antarctic icebergs. Von Suhm is said to have found three nests on high land near Christmas Harbour. I never met with any nests of this bird.]