

	Range in Depth. Fathoms.	Principal Localities	Tertiary.	Cretaceous.	Jurassic.
<i>Fibularia</i> , Lamk.					
× <i>F. australis</i> , Desm.	-950 C.	{ Sandwich Islands; Japan; Australia; Kingsmill Islands; New South Wales.			
<i>F. ovulum</i> , Launk.	—	{ Indian Ocean; Philippine Islands.	Eocene. Italy.	{ Damien. Belgium.	
× <i>F. volva</i> , Agass.	-28 C.	{ Red Sea; Formosa; North Australia.			
<i>ECHINANTHIDÆ</i> , A. Ag.					
<i>Clypeaster</i> , Lamk.					
× <i>C. humilis</i> , A. Ag.	-20 C.	{ Red Sea; East India Islands; New Caledonia; Philippines; Amboyna.			
<i>C. rotundus</i> , A. Ag.	—	{ Panama; San Diego.			
× <i>C. scutiformis</i> , Lamk.	—	{ Red Sea; Philippine Islands; Kingsmill Islands; Japan; New Caledonia.			
× <i>C. subdepressus</i> , Agass.	-120 B.	{ West Coast of Africa; South Carolina; Florida; Brazil; West India Islands.			
<i>Echinanthus</i> , Breyne.					
<i>E. rosaccus</i> , Gray.	-118 B.	{ West India Islands; Florida; South Carolina.			
× <i>E. testudinarius</i> , Gray.	-120 C.	{ Australia; Japan; Sandwich Islands; Gulf of California; New South Wales.			
<i>LAGANIDÆ</i> , Des. (Emend.)					
<i>Laganum</i> , Kl.					
<i>L. bonani</i> , Kl.	—	{ Tasmania; East India Islands; Philippine Islands.			
× <i>L. depressum</i> , Less.	—	{ Kingsmill Islands; Fiji Islands; Philippine Islands; Australia; Zanzibar; Arafura Sea.			
× <i>L. putnami</i> , Barn.	-25 C.	{ Japan; Amboyna.			
<i>Peronella</i> , Gray.					
× <i>P. decagonalis</i> , A. Ag.	-315 C.	{ Japan; New Caledonia; Bay of Bengal; East India Islands; Tahiti; Philippines; Tonga; Arafura Sea.			
<i>P. orbicularis</i> , A. Ag.	—	{ Australia; Formosa.			
× <i>P. peronii</i> , Gray.	-40 C.	{ Tasmania; Philippine Islands; New South Wales.			
<i>P. rostrata</i> , A. Ag.	—	New Zealand; Zanzibar.			
<i>SCUTELLIDÆ</i> , Agass.					
<i>Echinorachnius</i> , Leske.					
<i>E. excentricus</i> , Val.	—	{ California; Alaska; Kamtchatka.			
<i>E. mirabilis</i> , A. Ag.	—	{ Japan; Aleutian Islands.			
				North America; Patagonia; Australia.	