

	Range in Depth. Fathoms.	Principal Localities.
<i>Echinus</i> , Rond. (Linn.)		
× <i>E. acutus</i> , Lamk.,	-1350 C.	{ Norway; Shetland; Mediterranean; Halifax to New York; Ascension; Kermadec Islands.
<i>E. albocinctus</i> , Hutt.,	—	New Zealand.
× <i>E. angulosus</i> , A. Ag.,	—	{ Cape of Good Hope; Mauritius; Red Sea; Philippines; New Zealand; Australia.
<i>E. darleyensis</i> , Wood.,	—	Australia.
<i>E. elevatus</i> , Hutt.,	—	New Zealand.
<i>E. esculentus</i> , Linn.,	—	Norway; English Channel.
× <i>E. magellanicus</i> , Phil.,	-1600 C.	{ Patagonia; Chili; New Zealand; Australia; Cape of Good Hope to Marion Islands to Kerguelen; Falkland Islands.
× <i>E. margaritaceus</i> , Lamk.,	-245 C.	{ West Patagonia; Straits of Magellan; New Zealand; Ker- guelen; Heard Islands.
<i>E. melo</i> , Lamk.,	-190.	Mediterranean; Canary Islands; Cape Verde Islands.
<i>E. microtuberculatus</i> , Bl.,	—	Mediterranean; Cape Verde Islands.
<i>E. miliaris</i> , Müll.,	—	Norway; English Channel.
<i>Toxopneustes</i> , Agass.		
<i>T. maculatus</i> , A. Ag.,	—	Christmas Islands; Bourbon.
× <i>T. pileolus</i> , Agass.,	-10 C.	{ Panama; East India Islands; Fiji Islands; Mauritius; New Caledonia; Sandwich Islands; Philippines.
<i>T. semituberculatus</i> , Agass.,	—	Galapagos; West Coast Central America; Cape St Lucas.
× <i>T. variegatus</i> , A. Ag.,	-242 B.	{ Bermudas; South Carolina; West India Islands; Brazil; Straits of Florida.
<i>Hipponocē</i> , Gray.		
<i>H. depressa</i> , A. Ag.,	—	Gulf of California.
<i>H. esculenta</i> , A. Ag.,	-451 B.	Florida; West Indies; Surinam; Bermudas; Yucatan.
× <i>H. variegata</i> , A. Ag.,	-15 C.	{ Sandwich Islands; Japan; East India Islands; Fiji Islands; Red Sea; Mozambique; Philippines; Kandavu Reef; New South Wales.
<i>Evechinus</i> , Verrill.		
× <i>E. chloroticus</i> , Verrill.,	—	New Zealand; Kandavu Reef.
CLYPEASTRIDÆ, Agass.		
EUCLYPEASTRIDÆ, Haeckel.		
FIBULARINA, Gray.		
<i>Echinocyamus</i> , Van Phels.		
× <i>E. pusillus</i> , Gray,	-805 B.	{ Norway; Iceland; Mediterranean; Azores; Florida; Carib- bean Islands; North Brazil.
<i>Fibularia</i> , Lamk.		
× <i>F. australis</i> , Desml.,	-950 C.	{ Sandwich Islands; Japan; Australia; Kingsmills Islands; New South Wales.
<i>F. ovulum</i> , Lamk.,	—	Indian Ocean; Philippine Islands.
× <i>F. volva</i> , Agass.,	-28 C.	Red Sea; Formosa; North Australia.
ECHINANTHIDÆ, A. Ag.		
<i>Clypeaster</i> , Lamk.		
× <i>C. humilis</i> , A. Ag.,	-20 C.	{ Red Sea; East India Islands; New Caledonia; Philip- pines; Amboyna.
<i>C. rotundus</i> , A. Ag.,	—	Panama; San Diego.
× <i>C. scutiformis</i> , Lamk.,	—	{ Red Sea; Philippine Islands; Kingsmills Islands; Japan; New Caledonia.
× <i>C. subdepressus</i> , Agass.,	-120 B.	{ West Coast of Africa; Florida; South Carolina; Brazil; West India Islands.
<i>Echinanthus</i> , Breyn.		
<i>E. rosaceus</i> , Gray,	-118 B.	West India Islands; Florida; South Carolina.
× <i>E. testudinarius</i> , Gray,	-120 C.	{ Australia; Japan; Sandwich Islands; Gulf of California; New South Wales.