

Depth in Fathoms.	Name.	Geographical Distribution.
	<i>Parazetes</i> , Slater (1879).	
	Mandibles rudimentary, two-jointed ; palpi nine-jointed ; ovigerous legs ten-jointed ; proboscis fusiform.	
(?)	<i>Parazetes auchenicus</i> , Slater, Annals and Mag. of Nat. Hist., 5th series, iii. 281, 1879. Slater calls the generic characteristic of <i>Parazetes</i> "distinct." The description of the genus and species has been given from a single specimen, so I doubt whether the generic differences from <i>Zetes</i> or any other closely-allied genus will hold good.	Off Cape Sima (Japan).
	<i>Pariboea</i> , Philippi (1843).	
	Mandibles rudimentary, two-jointed ; palpi five-jointed ; ovigerous legs nine (none ten- (?)) jointed ; proboscis ovate.	
(?)	<i>Pariboea spinipalpis</i> , Philippi, Arch. f. Naturg., 1843, ix. p. 178. Perhaps the description of this species is not taken from an adult specimen.	Sorrento (Gulf of Naples).
	<i>Alcinous</i> , Costa (1861).	
	Mandibles rudimentary, three-jointed ; palpi seven- or nine-jointed ; ovigerous legs eight-jointed ; proboscis long-ovate.	
(?)	<i>Alcinous vulgaris</i> , Costa, Microdoride mediterranea, Napoli, 1861, p. 13. This species has, I believe, a certain resemblance to an <i>Ascorhynchus</i> .	Mediterranean.
(?)	<i>Alcinous megacephalus</i> , Costa, Microd. medit. Napoli, 1861, p. 14. This species belongs rather to <i>Pariboea</i> , Philippi, than to <i>Alcinous</i> , Costa.	Gulf of Naples.
	<i>Achelia</i> , Hodge (1864).	
	Mandibles rudimentary, two-jointed ; palpi eight-jointed ; ovigerous legs ten-jointed ; proboscis pyriform.	
Low water mark to 17 fathoms; 35 fathoms.	<i>Achelia spinosa</i> , Stimpson (sp.). <i>Zetes spinosa</i> , Stimpson. Invertebrata of Grand Manan, 1853, p. 37. <i>A. spinosa</i> , Wilson, Pycnogon. of New England, Trans. Connect. Acad. of Arts and Sciences, vol. v. p. 7, 1880. The differences of <i>A. spinosa</i> , Stimpson, from Hodge's <i>A. echinata</i> "seem scarcely sufficient to separate them as distinct species" (Wilson, <i>loc. cit.</i>).	Grand Manan ; Eastport, Casco Bay, off Block Island, &c.
Low tide to a few faths. (Hodge).	<i>Achelia echinata</i> , Hodge, British Pycnogonoidea, Ann. and Mag. of Nat. Hist., 3d ser., vol. xiii. p. 115, 1864. A good description of this species has been given by Grube, St Vaast la Hougue und seine Meeres—besonders seine Anneliden-Fauna, Verhandl. d. Schles. Gesellsch. f. vaterl. Cultur, 1869-72, p. 27 (Separatabdruck). Perhaps identical with <i>A. spinosa</i> , Stimpson, Invertebrata of Grand Manan, 1853, p. 37.	Channel Islands, Isle of Man ; Durham Coast, St Vaast la Hougue, Dutch Coast.