

t T. TENTACULA.

TENTACLES.

TENTAKELN.

The corona of tentacles in the Medusæ corresponds originally to that of the nearly related Polyps, and is therefore placed on the margin of the umbrella in the Medusæ, as it is on the margin of the peristome in the Polyps; in many Medusæ, however, the tentacles assume a secondary position, and are sometimes transferred to the dorsal, sometimes to the ventral surface of the umbrella.

<i>ta</i>	Tentacula adradialia.	Adradial tentacles.	Adradiale Tentakeln.
<i>tb</i>	Tentaculorum bulbi basales.	Basal bulbs of the tentacles.	Basale Tentakel-Bulben.
<i>tc</i>	Tentacula cava.	Hollow tentacles.	Hohle Tentakeln.
<i>td</i>	Tentacula solida.	Solid tentacles.	Dichte Tentakeln.
<i>te</i>	Cirri spirales.	Spiral cirri.	Spiral-Fäden.
<i>ti</i>	Tentacula interradialia.	Interradial tentacles.	Interradiale Tentakeln.
<i>tk</i>	Tentacula ancoralia.	Margin anchors or marginal papillæ.	Randanker oder Randpapillen.
<i>tp</i>	Tentacula perradialia.	Perradial tentacles.	Perradiale Tentakeln.
<i>tr</i>	Tentaculorum radices.	Roots of the tentacles.	Tentakel-Wurzeln.
<i>ts</i>	Tentacula suctoria.	Suckers.	Saugnäpfe.

U. UMBRELLA.

UMBRELLA.

SCHIRM.

<i>u</i>	<i>Discus, campana,</i> <i>Discus gelatinosus.</i>	<i>Disc, bell,</i> <i>Gelatinous disk.</i>	<i>Scheibe, Glocke,</i> <i>Gallertscheibe.</i>
----------	-------------------------------------------------------	-----------------------------------------------	---------------------------------------------------

The gelatinous concavo-convex disk, the most voluminous part of the Medusa, and in most cases forming the principal part of the body; the upper convex surface (exumbrella) passes into the lower concave surface (subumbrella), at the margin of the umbrella.

<i>ua</i>	Umbrellæ pedalia adradialia.	Adradial pedalia.	Adradiale Schirm-Pedale.
<i>uc</i>	Umbrellæ conus (cupula).	Cone of the umbrella.	Schirmkegel (Schirmkuppel).
<i>ud</i>	Umbrellæ pedalia.	Pedalia of the umbrella.	Schirm-Pedale.
<i>uf</i>	Umbrellæ fibrillæ elasticæ.	Elastic fibrillæ of the gelatinous disk.	Elastische Fasern der Schirm-Gallerte.
<i>ug</i>	Umbrellæ gelatina.	Gelatinous substance of the umbrella.	Schirm-Gallerte.
<i>ui</i>	Umbrellæ pedalia interradialia.	Interradial pedalia.	Interradiale Schirm-Pedale.
<i>uk</i>	Umbrellæ conus lingualis.	Tongue-like cone of the umbrella.	Zungenkegel oder Gastralkegel.
<i>um</i>	Margo umbralis.	Margin of the umbrella.	Schirmrand.
<i>up</i>	Umbrellæ pedalia perradialia.	Perradial pedalia.	Perradiale Schirm-Pedale.
<i>us</i>	Pedunculus gastralis.	Gelatinous peduncle of the stomach.	Magenstiel (gallertiger, solider Stiel).
<i>ut</i>	Tubercula umbrellæ.	Tuberles of the umbrella.	Schirmhöker.
<i>uz</i>	Umbrellæ corona.	Corona of the umbrella.	Peripherer Schirmkranz.

v V. VELUM.

VELUM.

RANDHAUT.

A muscular, thin, membranous ring, which, in the section of the Craspedota, projects freely downwards and inwards from the umbrella margin, as a process of the subum-