

160; from the South Pacific Ocean, Stations 281 and 299, and from the North Pacific Ocean, Stations 241 and 244.

GEOGRAPHICAL DISTRIBUTION OF THE ELASIPODA.

NORTH ATLANTIC OCEAN.

STATION V.—Lat. $35^{\circ} 47' N.$, long. $8^{\circ} 23' W.$ Depth, 1090 fms. Mud.

Benthodytes typica, n. sp.

Euphronides depressa, n. sp.

STATION 23.—Off Sombrero Island. Depth, 450 fms. Globigerina ooze.

Orphnurgus asper, Théel.

STATION 50.—Lat. $42^{\circ} 8' N.$, long. $63^{\circ} 39' N.$ Depth, 1250 fms. Grey ooze.

Kolga nana, Théel.

STATION 101.—Lat. $5^{\circ} 48' N.$, long. $14^{\circ} 20' W.$ Depth, 2500 fms. Mud.

Psychropotes semperiana, n. sp.

STATION 104.—Lat. $2^{\circ} 25' N.$, long. $20^{\circ} 1' W.$ Depth, 2500 fms. Grey ooze.

Peniagone lugubris, n. sp.

SOUTH ATLANTIC OCEAN.

STATION 133.—Lat. $35^{\circ} 41' S.$, long. $20^{\circ} 55' W.$ Depth, 1900 fms. Globigerina ooze.

Psychropotes semperiana, n. sp.

STATION 325.—Lat. $36^{\circ} 44' S.$, long. $46^{\circ} 16' W.$ Depth, 2650 fms. Grey mud.

Scotoplanes papillosa, n. sp.

Oneirophanta mutabilis, Théel.

SOUTH INDIAN OCEAN.

STATION 143.—Lat. $36^{\circ} 48' S.$, long. $19^{\circ} 24' E.$ Depth, 1900 fms. Globigerina ooze.

Scotoplanes albida, n. sp.

STATION 146.—Lat. $46^{\circ} 46' S.$, long. $45^{\circ} 31' E.$ Depth, 1375 fms. Globigerina ooze.

Psychropotes lovéni, n. sp.

Oneirophanta mutabilis, Théel.