

Parelpidia elongata, Théel.

- Fig. 16. Four-armed deposits from the body-wall; the arms about 0.12 mm. long.
 „ 17. Spicula from the pedicels.

Elpidia rigida, n. sp.

- „ 18. Four-armed deposits from the body-wall; the arms about 0.48 mm. long.
 „ 19. Spicula and four-armed deposits from the pedicels.
 „ 20. Deposits from the tentacles.

Achlyonice lactea, n. sp.

- „ 21. Three-armed deposits with the longest arms about 0.22 mm.; and a small wheel measuring 0.06 mm. in diameter. Taken from the dorsal perisoma.
 „ 22. Deposits from the ventral perisoma.
 „ 23. Spicula from the pedicels.

PLATE XXXIII.

Kolga nana, Théel.

- Fig. 1. Deposits from the body-wall, about 0.08 mm. long.
 „ 2. Spicula from the pedicels.

Elpidia incerta, n. sp.

- „ 3. Four-armed deposits from the body-wall, with the arms from 0.06 mm. to 0.16 mm.
 „ 4. Deposits from the pedicels.

Peniagone atrox, n. sp.

- „ 5. Larger and smaller four-armed deposits from the body-wall; the arms of the largest deposits 0.12 mm. long.

Elpidia ambigua, n. sp.

- „ 6. Four-armed deposit with the arms 0.18 mm. long, and wheel, measuring 0.048 mm. in diameter.

Scotoplanes insignis, n. sp.

- „ 7. Three-armed deposit and C-shaped bodies; the arms of the former 0.24 mm. long, and the latter measuring about 0.1 mm. in length.

Peniagone horrifer, n. sp.

- „ 8. Deposits from the pedicels.
 „ 9. Four-armed deposits from the body-wall, with the arms about 0.06 mm. long.