

Species of *Pectinura* not herein described.

Pectinura vestita, Fbs., Linn. Trans., vol. xix. p. 143, pl. xiii. figs. 1-7, 1842; Ltk., Addit. ad Hist., vol. iii. p. 31; Lym., Bull. Mus. Comp. Zool., vol. iii., part 10, p. 222; Ludwig, Echin. des Mittelmeeres, p. 546.

Ophiarachna vestita, Lym., Ill. Cat. Mus. Comp. Zool., vol. i. p. 11.

Ægean Sea; South Adriatic; 100 fathoms.

Pectinura infernalis, Ltk., Addit. ad Hist., part 3, p. 33, 1869; Lym., Bull. Mus. Comp. Zool., vol. iii. part 10, p. 222, pl. vii. fig. 1.

Ophiarachna infernalis, Müll. & Tr., Syst. Ast., p. 105, 1842.

Great Ocean.

Pectinura marmorata, Lym., Bull. Mus. Comp. Zool., vol. iii., part 10, p. 222, pl. v. figs. 1-7, 1874.

Philippines.

Pectinura septemspinosa, Ltk., Addit. ad Hist., part 3, p. 33, 1869; Lym., Bull. Mus. Comp. Zool., vol. iii., part 10, p. 222, pl. vi. figs. 10-13.

Ophiura septemspinosa, Kuhl. & v. Has., MS.

Ophiarachna septemspinosa, Müll. & Tr., Syst. Ast., p. 105, 1842.

Great Ocean.

Pectinura spinosa, Lym., Bull. Mus. Comp. Zool., vol. iii., part 10, p. 221, 1874.

Ophiarachna spinosa, Ljn., Oph. Viv. Öf Kong. Akad., p. 305, 1866.

Ophiopezella spinosa, Ljn., Dr. Goës, Oph. Öf. Kong. Akad., p. 639, 1871.

Foua Islands.

Pectinura verrucosa, Studer, Monatsb. König. Akad. Berlin, p. 461, July 1876.
Kerguelen Island; 150 fathoms.

Ophiopæpale.

Ophiopæpale, Ljn., Dr. Goës, Oph. Öf. Kong. Akad., 1871.

Disk granulated. Teeth, and numerous even, close-set, mouth papillæ. No tooth papillæ. Three slender, short, smooth arm spines. One small tentacle scale. Arms long, cylindrical, and gradually tapering to a fine point, their under plates divided into two parts, an inner tongue, and an outer piece which is of a transverse oblong shape. Two genital openings in each interbrachial space.