

10 mm. six to seven spines. The figure of the original *Amphiura chiajei* (Fbs. *loc. cit.*) agrees well with this northern form, except in having a large central rosette of primary plates.

*Amphiura eugeniae*, Ljn., Oph. Viv. Öf. Kong. Akad., p. 318, 1866 ; Dr. Goës, Oph. Öf. Kong. Akad., p. 642, 1871 ; Lym., Ill. Cat. Mus. Comp. Zool., No. viii., part 2, p. 5.

Atlantic, off Rio La Plata ; 30 to 55 fathoms.

*Amphiura palmeri*, Lym.

*Amphiura flexuosa* (?), Lym., Ill. Cat. Mus. Comp. Zool., No. viii., part 2, p. 17, pl. iii. figs. 35-37, pl. v. fig. 68, 1875.

West Indies ; 100 fathoms.

*Amphiura magellanica*, Ljn., Oph. Viv. Öf. Kong. Akad., p. 320, 1866 ; Lym., Ill. Cat. Mus. Comp. Zool. (same sp. ?). No. viii., part 2, p. 19.

Straits of Magellan ; North-east Patagonia ; 30 fathoms.

*Amphiura grandisquama*, Lym., Bull. Mus. Comp. Zool., vol. i., part 10, p. 334, 1869 ; Bull. Mus. Comp., vol. v., part 9, p. 226.

West Indies ; 10 to 240 fathoms.

*Amphiura stimpsoni*, Ltk., Addit. ad Hist., part 2, p. 116, 1859.

West Indies ; Cape Frio, Brazil ; 10 to 35 fathoms.

*Amphiura duncani*, Lym.

*Amphiura Lütkeni*, Duncan, Journ. Linn. Soc., vol. xiv., p. 464, pl. x. fig. 17, 1879.

Korean Sea.

*Amphiura sundevalli*, Ljn., Oph. Viv. Öf. Kong. Akad., p. 320, 1866.

*Ophiolepis Sundevalli*, Müll. & Tr., Syst. Ast., p. 93, 1842.

*Amphiura Holbölli*, Ltk., Vid. Meddel., Nov. 1854, p. 98 ; Addit. ad Hist., part 1, p. 55, pl. ii. figs. 13 a, b ; Lym., Ill. Cat. Mus. Comp. Zool. No. i. p. 118.

European Arctic Sea ; Greenland ; 15 to 50 fathoms.

*Amphiura semiermis*, Lym., Bull. Mus. Comp. Zool., vol. i., part 10, p. 332, 1869 ; Bull. Mus. Comp. Zool., vol. v., part 9, p. 227.

West Indies ; 377 to 539 fathoms.