

<i>Ophiomyxa flaccida</i> , I., II.	<i>Astroschema oligactes</i> , II.
<i>australis</i> , I., II.	<i>brachiatum</i> .
<i>vivipara</i> , II.	<i>rubrum</i> .
<i>Ophiochondrus convolutus</i> .	<i>sulcatum</i> .
<i>Hemieuryale pustulata</i> , II.	<i>intectum</i> .
<i>Sigsbeia murrhina</i> , II.	<i>Ophiocreas lumbricus</i> , II.
<i>Gorgonocephalus malmgrenii</i> .	<i>carnosus</i> .
<i>Astrogomphus vallatus</i> , II.	<i>caudatus</i> .
<i>Astrochele lymani</i> , IV.	<i>ædipus</i> .
<i>Astrotoma murrayi</i> .	<i>Astronyx loveni</i> , II.

Number of Species in Table III., 137.

TABLE IV.—Species appearing below 500 and above 1000 fathoms.

<i>Pectinura heros</i> .	<i>Ophiactis poa</i> , V.
<i>Ophiozona</i> (?) <i>dubia</i> .	<i>Amphiura bellis</i> , III.
<i>Ophioceramis</i> (?) <i>clausa</i> .	<i>otteri</i> , V.
<i>Ophiopleura borealis</i> .	<i>argentea</i> .
<i>artica</i> (no depth given but probably belongs here).	<i>acacia</i> .
<i>Ophioglypha meridionalis</i> , V.	<i>iris</i> , III.
<i>palliata</i> , III., V.	<i>semiermis</i> , III.
<i>lepida</i> , III., V.	<i>lanceolata</i> .
<i>irrorata</i> , III., V.	<i>sarsii</i> , I.—III.
<i>albata</i> .	<i>lunaris</i> .
<i>variabilis</i> , III., V.	<i>duplicata</i> , II., III., V.
<i>lapidaria</i> .	<i>concolor</i> , V.
<i>rugosa</i> .	<i>canescens</i> .
<i>confragosa</i> .	<i>Amphilepis norvegica</i> , II., III., V.
<i>Ophiomusium planum</i> .	<i>Ophiochiton lentus</i> .
<i>lymani</i> , V.	<i>Ophiacantha vepratrica</i> .
<i>validum</i> , III., V.	<i>rosea</i> , II., III.
<i>scalare</i> .	<i>vivipara</i> , I.—III.
<i>pulchellum</i> , III., V.	<i>bidentata</i> , II., III., V.
<i>Ophiomastus tegulitius</i> , III., V.	<i>stimulea</i> .
<i>Ophiophyllum petilum</i> .	<i>echinulata</i> .
<i>Ophiopholis japonica</i> , III.	<i>abnormis</i> , III.
<i>Ophiactis flexuosa</i> , III.	<i>smittii</i> .
<i>cuspidata</i> .	<i>cornuta</i> .
<i>nama</i> , III.	<i>cosmica</i> , III., V.
	<i>pentacrinus</i> , II., III.