

TEMPERATURE TABLES.

TABLE I.—Showing Challenger species found between 32° and 38° Fahr. inclusive.

The figure after each species indicates its depth in fathoms. The Roman numeral opposite a species shows that it is found also in one of the other tables thus indicated.

<i>Ophiozona stellata</i> , 1100, II.
<i>Ophioplithus medusa</i> , ¹ 1975.
<i>grisea</i> , ¹ 1975.
<i>Ophiernus vallincola</i> , 1000–175.
<i>Ophioglypha sarsii</i> , 3–238, II.
<i>meridionalis</i> , 600–1035.
<i>levida</i> , 425–1350, II.
<i>æqualis</i> , 1070.
<i>irrorata</i> , 410–1900.
<i>orbiculata</i> , 1875.
<i>undulata</i> , 1850.
<i>albata</i> , 775.
<i>loveni</i> , 1375–2600.
<i>fraterna</i> , 1950.
<i>bullata</i> , 1240–2850.
<i>convexa</i> , 2050–2350.
<i>sculptilis</i> , 1875.
<i>variabilis</i> , 390–1425.
<i>ornata</i> , 2160–2600.
<i>lacazei</i> , 2160–2600.
<i>lienosa</i> , 1950.
<i>radiata</i> , 1050.
<i>undata</i> , 1450.
<i>lapidaria</i> , 565.
<i>minuta</i> , 1875–1800.
<i>deshayesi</i> , 28–150.
<i>inornata</i> , 1850.
<i>confragosa</i> , 600.
<i>Ophiocten amitimum</i> , 120–1260.
<i>umbraticum</i> , 2650.
<i>pallidum</i> , 2600.
<i>hastatum</i> , 1375.

<i>Ophiomusium armigerum</i> , 1650–2200.
<i>corticosum</i> , 1850.
<i>cancellatum</i> , 420–470.
<i>archaster</i> , 1900.
<i>lymani</i> , 565–1825.
<i>granosum</i> , 1875.
<i>Ophiomastus tegulitius</i> , 275–2600, II.
<i>Ophioconis antarctica</i> , 50–150.
<i>Ophiodolos aculeata</i> , littoral to 400, II.
<i>japonica</i> , 420–775.
<i>Amphiura bellis</i> , 210–775, II.
<i>otteri</i> (?), 550–1250, II.
<i>studerii</i> , 20–310.
<i>acacia</i> , 565.
<i>iris</i> , 420–775.
<i>angularis</i> , 150.
<i>squamata</i> , littoral to 120,
II., III.
<i>duplicata</i> , 78–1560.
<i>concolor</i> , 800–1425, II.
<i>dalea</i> , 2650.
<i>cernua</i> , 2300.
<i>glaucha</i> , 345–420, II.
<i>Amphilepis norvegica</i> , 50–1350.
<i>patens</i> , 2160.
<i>tenuis</i> , 1875.
<i>Ophiocymbium cavernosum</i> , 1950.
<i>Ophiochytra epigrus</i> , 2350.
<i>Ophiacantha placentigera</i> , 1350.
<i>rosea</i> , 150–775.
<i>vivipara</i> , 20–600, II.
<i>bidentata</i> , 5–1350.

¹ No temperature given, but at 1975 fathoms; it must fall within the limits of this table.