

Station 192, September 26, 1874; off Ki Island, lat.  $5^{\circ} 42' S.$ , long.  $132^{\circ} 25' E.$ ;  
Noon; depth, 129 fathoms; bottom, mud.

*Polycarpa quadrata.*

*Ecteinascidia crassa.*

Banda, Malay Archipelago, October 1, 1874; depth, 17 fathoms.

*Polycarpa sulcata.*

*Ecteinascidia fusca.*

Between Hong Kong and New Guinea the following Tunicata were collected:—

Hong Kong, China; depth, 10 fathoms.

*Corella japonica.*

Station 208, January 17, 1875; lat.  $11^{\circ} 37' N.$ , long.  $123^{\circ} 32' E.$ ; depth, 18 fathoms; bottom, mud.

*Polycarpa irregularis.*

Station 212, January 30, 1875; lat.  $6^{\circ} 55' N.$ , long.  $122^{\circ} 15' E.$ ; depth, 10, 14 and 20 fathoms; bottom, sand.

*Polycarpa pedata.*

Between New Guinea and Japan the following Tunicata were dredged:—

Station 233A, May 17 to 19, 1875; Kobé, Japan; lat.  $34^{\circ} 35' N.$ , long.  $135^{\circ} 10' E.$ ; depth, 8 fathoms and 50 fathoms; bottom, mud and sand.

*Styela clava.*

*Corella japonica.*

*Ciona savignyi.*

Off Yokohama, Japan; shallow water.

*Corella japonica.*

Between Japan and the Sandwich Islands the following Tunicata were obtained:—

Station 241, June 23, 1875; lat.  $35^{\circ} 41' N.$ , long.  $157^{\circ} 42' E.$ ; depth, 2300 fathoms; bottom temperature,  $1^{\circ} 1 C.$ ; bottom, red clay.

*Culeolus murrayi.*

Station 248, July 5, 1875; lat.  $37^{\circ} 41' N.$ , long.  $177^{\circ} 4' W.$ ; depth, 2900 fathoms; bottom temperature,  $1^{\circ} 1 C.$ ; bottom, red clay.

*Hypobythius calycodes.*

Between the Sandwich Islands and Valparaiso the following Tunicata were collected:—

Papiete Harbour, Tahiti, Society Islands; depth, 10 to 20 fathoms.

*Cynthia papietensis.*