

Specimen.	Length of left hepatic duct.	Distance from pylorus of point of entrance into intestine of left hepatic duct.	Length of right hepatic duct.	Distance from pylorus of point of entrance into intestine of right hepatic duct.	Length of gall bladder.
<i>Eudyptes chrysocome</i> , from Kerguelen.					
No. 1	$3\frac{3}{4}$	4	4	11	$3\frac{1}{2}$
No. 2	$2\frac{3}{4}$	$3\frac{3}{4}$	3	8	3
<i>Eudyptes chrysolophus</i> .					
No. 1	3	13	4	9	$4\frac{1}{2}$
No. 2	3	3	4	8	$4\frac{1}{2}$
<i>Spheniscus demersus</i> .					
No. 1	$3\frac{1}{2}$	14	4	21	4
No. 2	$3\frac{3}{4}$	9	$4\frac{1}{4}$	15	...
No. 3 immature.	...	8	...	$11\frac{1}{2}$...
<i>Spheniscus magellanicus</i> .					
No. 1	5	13	7	26	$5\frac{1}{2}$
No. 2	5	13	$5\frac{1}{2}$	25	...
<i>Spheniscus mendiculus</i> .					
No. 1	3	$5\frac{1}{4}$	3	$7\frac{1}{2}$	3
<i>Spheniscus minor</i> .					
No. 1	...	$2\frac{1}{2}$...	$4\frac{3}{4}$	$2\frac{1}{4}$