

PLATE VII.

OSTEOLOGY OF PENGUINS.

- Fig. 1.—Right scapula of *Eudyptes chrysocome*, from Tristan d'Acunha ; natural size.
- Fig. 2.—Right scapula of *Spheniscus demersus* ; natural size.
- Fig. 3.—Right scapula of *Pygosceles tæniatus* ; two-thirds natural size.
- Fig. 4.—Right scapula of *Aptenodytes longirostris* ; two-thirds natural size.
- Fig. 5.—Left coracoid bone of *Eudyptes chrysocome* ; natural size.
- Fig. 6.—Left coracoid bone of *Pygosceles tæniatus* ; two-thirds natural size.
- Fig. 7.—Interior of pelvis of *Eudyptes chrysocome*, from Tristan d'Acunha ; natural size.
- Fig. 8.—Exterior of pelvis of *Eudyptes chrysocome*, from Tristan d'Acunha ; natural size.
- Fig. 9.—Patella of *Eudyptes chrysocome*, from Tristan d'Acunha, outer surface ; natural size.
- Fig. 10.—Patella of *Eudyptes chrysocome*, from Tristan d'Acunha, seen from the front ; natural size.
- Fig. 11.—Right tarso-metatarsal bone of *Eudyptes chrysocome*, from Tristan d'Acunha ; natural size.
- Fig. 12.—Right tarso-metatarsal bone of *Spheniscus demersus* ; natural size.
- Fig. 13.—Right tarso-metatarsal bone of *Pygosceles tæniatus* ; natural size.
- Fig. 14.—Right tarso-metatarsal bone of *Aptenodytes longirostris* ; natural size.