

Seguenza's fossil specimens were from the Miocene deposits of Sicily. It is also recorded in Wright's catalogue of the Post-tertiary Foraminifera of the north-east of Ireland.

*Lagena marginata*, Walker and Boys (Pl. LIX. figs. 21–23).

"Serpula ( <i>Lagena</i> ) marginata,"	Walker and Boys, 1784, Test. Min., p. 2, pl. i. fig. 7.
<i>Vermiculum marginatum</i> , Montagu, 1803, Test. Brit., p. 524.	
<i>Oolina compressa</i> , d'Orbigny, 1839, Foram. Amér. Mérid., p. 18, pl. v. figs. 1, 2.	
<i>Entosolenia marginata</i> (pars), Williamson, 1848, Ann. and Mag. Nat. Hist., ser. 2, vol. i. p. 17, pl. ii. figs. 15–17.	
<i>Fissurina alata</i> , Reuss, 1851, Zeitschr. d. deutsch. geol. Gesellsch., vol. iii. p. 58, pl. iii. fig. 1.	
,, <i>carinata</i> , Reuss, 1862, Sitzungsb. d. k. Ak. Wiss. Wien, vol. lxvi. p. 338, pl. vi. fig. 83; pl. vii. fig. 86.	
,, <i>communis</i> , Seguenza, 1862, Foram. Monat. Mess., p. 59, pl. i. figs. 56, 57.	
,, <i>propinqua</i> , Id.	Ibid. p. 59, pl. i. fig. 58.
,, <i>ardasii</i> , Id.	Ibid. p. 59, pl. i. fig. 59.
,, <i>aperta</i> , Id.	Ibid. p. 60, pl. i. fig. 60.
,, <i>obvia</i> , Id.	Ibid. p. 60, pl. ii. fig. 1.
,, <i>tenuis</i> , Id.	Ibid. p. 60, pl. ii. fig. 2.
,, <i>elliptica</i> , Id.	Ibid. p. 60, pl. ii. fig. 3.
,, <i>ovata</i> , Id.	Ibid. p. 62, pl. ii. figs. 9, 10.
,, <i>benoitiana</i> , Id.	Ibid. p. 62, pl. ii. fig. 11.
,, <i>haeckelii</i> , Id.	Ibid. p. 63, pl. ii. fig. 13.
,, <i>circulum</i> , Id.	Ibid. p. 63, pl. ii. fig. 15.
,, <i>silvestrii</i> , Id.	Ibid. p. 64, pl. ii. fig. 18.
,, <i>emarginata</i> , Id.	Ibid. p. 65, pl. ii. fig. 20.
,, <i>dilatata</i> , Id.	Ibid. p. 65, pl. ii. fig. 21.
,, <i>lævis</i> , Id.	Ibid. p. 66, pl. ii. figs. 22, 23.
,, <i>sulcata</i> , Id.	Ibid. p. 67, pl. ii. fig. 29.
,, <i>tubulosa</i> , Id.	Ibid. p. 68, pl. ii. figs. 36, 37.
,, <i>regolaris</i> , Id.	Ibid. p. 71, pl. ii. fig. 46.
,, <i>sartorii</i> , Id.	Ibid. p. 71, pl. ii. fig. 47.
,, <i>lyellii</i> , Id.	Ibid. p. 71, pl. ii. figs. 48, 49.
<i>Lagena sulcata</i> , var. ( <i>Entosolenia</i> ) <i>marginata</i> , Parker and Jones, 1865, Phil. Trans., vol. clv. p. 355, pl. xiii. figs. 42–44; pl. xvi. fig. 12, a. b.	
<i>Fissurina alata</i> , Reuss, 1870, Sitzungsb. d. k. Ak. Wiss. Wien, vol. lxii. p. 469, No. 2;—Schlicht, 1870, Foram. Pietzpuhl, pl. iv. figs. 7–9, 13–15; pl. v. figs. 19–21.	
<i>Lagena vulgaris</i> , var. <i>petasma-marginata</i> , Ry. Jones, 1872, Trans. Linn. Soc. Lond., vol. xxx.	p. 58, pl. xix. figs. 38, 39.
,, " var. <i>bracteato-marginata</i> , Id.	Ibid. p. 59, pl. xix. figs. 40, 41.
,, " var. <i>alato-marginata</i> , Id.	Ibid. p. 60, pl. xix. fig. 44.
,, " var. <i>clavato-marginata</i> , Id.	Ibid. p. 60, pl. xix. fig. 45.
,, " var. <i>pertuso-marginata</i> , Id.	Ibid. p. 61, pl. xix. figs. 46, 47.

*Lagena marginata* includes those compressed varieties of the type of which the peripheral edge is carinate to a greater or less degree. In some cases the carina is only sufficient to form a thin sharp margin, whilst in others it is developed into a lamelliform