

PLATE XXVI.

PELOSINA.—HIPPOCREPINA.—BATHYSIPHON.

	Diam.	Page
Figures 1-6.— <i>Pelosina cylindrica</i> , n. sp., Figs. 1-3. Lateral aspect ; fig. 4, oral end ; fig. 5, aboral end. Fig. 6. Specimen laid open to show the thickness of the walls and the smooth interior.	× 9	236
„ 7-9.— <i>Pelosina variabilis</i> , Brady, Figs. 7, 8. General aspect of the test. Fig. 9. Portion of a test laid open to show the thick- ness of the walls.	× 9	235
„ 10-14.— <i>Hippocrepina indivisa</i> , Parker, Figs. 10, 11. Lateral aspect ; figs. 12, 13, oral end. Fig. 14. Test laid open longitudinally, showing the undivided cavity.	× 45	325
„ 15-20.— <i>Bathysiphon filiformis</i> , M. Sars, Figs. 15, 16. General aspect of the test, Fig. 17. Portion of a specimen laid open longitudinally, to show the structure of the walls and the nature of the internal cavity, Fig. 18. Transverse section of the test, viewed by transmitted light, Fig. 19. Longitudinal section of the wall, with a mass of adherent dark-coloured sarcode, Fig. 20. Longitudinal section, more highly magnified,	× 3 × 60 × 60 × 60 × 200	248