

PLATE LXXXIV.

SPHÆROIDINA.—PULLENIA.

	Diam.	Page
<p>Figures 1-7.—<i>Sphæroidina bulloides</i>, d'Orbigny, . . .</p> <p style="padding-left: 40px;">Figs. 1-4, 6, 7. Specimens of various forms, in different aspects.</p> <p style="padding-left: 40px;">Fig. 5. Section of the shell, showing the laminated condition of the walls.</p>	× 50	620
<p>„ 8-11.—<i>Sphæroidina dehiscens</i>, Parker and Jones, . . .</p> <p style="padding-left: 40px;">Figs. 8, 9. Bottom specimens.</p> <p style="padding-left: 40px;">Fig. 10. Surface specimen.</p> <p style="padding-left: 40px;">Fig. 11. Section of the shell, showing the thickness of the walls and their coarse tubulation.</p>	× 50	621
<p>„ 12, 13.—<i>Pullenia sphaeroides</i>, d'Orbigny, sp., . . .</p> <p style="padding-left: 40px;"><i>a, a</i>, Lateral aspect; <i>b, b</i>, peripheral aspect.</p>	× 50	615
<p>„ 14, 15.—<i>Pullenia quinqueloba</i>, Reuss,</p> <p style="padding-left: 40px;">Figs. 14 <i>a</i>, 15. Lateral aspect; fig. 14 <i>b</i>, peripheral aspect.</p>	× 50	617
<p>„ 16-20.—<i>Pullenia obliquiloculata</i>, Parker and Jones, . . .</p> <p style="padding-left: 40px;">Figs. 16, 17. Bottom specimens. <i>a, a</i>, Lateral aspect <i>b, b</i>, peripheral aspect.</p> <p style="padding-left: 40px;">Fig. 18. Surface specimen.</p> <p style="padding-left: 40px;">Fig. 19. Monstrous example, isomorphous with <i>Globigerina helicina</i>.</p> <p style="padding-left: 40px;">Fig. 20. Section of the shell, showing the thickness and tubulation of the walls, and the infolding of the superior margin of the septal apertures.</p>	× 50	618