

GEOGRAPHICAL DISTRIBUTION OF THE COPEPODA COLLECTED DURING THE CRUISE
OF H.M.S. CHALLENGER.

	1	2	3	4	5	6	7
	North Atlantic.	South Atlantic.	South Indian Ocean.	Austral- asia.	South Pacific.	North Pacific.	Eastern Asia.
CALANIDÆ.							
<i>Calanus finmarchicus</i> (Günner),				+	+		
<i>propinquus</i> , n. sp.,	+	+	+	+	+	+	
<i>valgus</i> , n. sp.,	+	+		+	+		
<i>tonsus</i> , n. sp.,		+		+	+		
<i>gracilis</i> , Dana,	+	+		+	+		+
<i>princeps</i> , n. sp.,	+						
<i>Eucalanus attenuatus</i> , Dana,	+		+	+	+	+	+
<i>setiger</i> , n. sp.,	+			+			+
<i>Rhincalanus cornutus</i> , Dana,	+						+
<i>gigas</i> , n. sp.,		+	+			+	+
<i>Hemicalanus longicornis</i> , Claus,	+	+			+		
<i>orientalis</i> , n. sp.,				+			
<i>aculeatus</i> , n. sp.,			+			+	
<i>Pleuromma abdominale</i> (Lubbock),	+	+	+	+	+	+	
<i>Heterochæta spinifrons</i> , Claus,	+	+	+		+		
<i>Leuckartia flavicornis</i> , Claus,	+	+	+	+	+		
<i>scopularis</i> , n. sp.,						+	
<i>Undina vulgaris</i> Dana,	+	+		+		+	+
<i>darwinii</i> , Lubbock,	+	+		+	+	+	+
<i>Scolecithrix danæ</i> (Lubbock),	+	+		+	+	+	+
<i>minor</i> , n. sp.,			+				
<i>Euchæta prestandreæ</i> , Philippi,	+	+	+	+	+	+	+
<i>hessei</i> , n. sp.,	+			+	+	+	+
<i>pulchra</i> (Lubbock),	+						
<i>philippii</i> , n. sp.		+			+		
<i>australis</i> , n. sp.,		+		+			
<i>gigas</i> , n. sp.,		+					
<i>barbata</i> , n. sp.,		+					
<i>Calanoides patagoniensis</i> , n. gen. and sp.,					+		
<i>Ætidius armatus</i> , n. gen. and sp.,	+	+	+	+			+
<i>Drepanopus pectinatus</i> , n. gen. and sp.,			+				
<i>furcatus</i> , n. gen. and sp.,	+			+	+		
<i>Phyllopus bidentatus</i> , n. gen. and sp.,		+					
<i>Temora dubia</i> (Lubbock),	+	+		+	+		+
<i>armata</i> , Claus,	+						
<i>Centropages brachiatus</i> (Dana),					+		
<i>furcatus</i> (Dana),				+			+
<i>violaceus</i> (Claus),	+	+		+	+	+	+
<i>Candace pectinatu</i> , Brady,	+			+			+
<i>pachydactyla</i> , Dana,	+	+		+	+		+
<i>truncata</i> , Dana,	+	+	+	+		+	+