

LIST OF THE PRINCIPAL LOCALITIES FROM WHICH COPEPODA WERE
TAKEN, WITH THE NAMES OF THE SPECIES FOUND IN EACH.

Except where otherwise stated, all these gatherings were made during the daytime by means of the surface-net.

STATION 45.—May 3, 1873. Lat. $38^{\circ} 34'$ N., long. $72^{\circ} 10'$ W. Depth 1240 fathoms.

Bottom temperature, $2^{\circ} 4$ C. Mud (dredge).

Calanus princeps, n. sp.

STATION 50.—May 21, 1873. Lat. $42^{\circ} 8'$ N., long $63^{\circ} 39'$ W. Depth 1250 fathoms.

Bottom temperature, $2^{\circ} 8$ C. Grey ooze (dredge).

Calanus Princeps, n. sp.

May 22, 1873. Lat. $41^{\circ} 19'$ N., long. $63^{\circ} 12'$ W. South of Halifax,
Nova Scotia.

Saphirina opaca, Lubbock.

STATION 99.—August 15, 1873. Lat. $7^{\circ} 53'$ N., long. $17^{\circ} 26'$ W.; also August 17, 1873.

Miracia efferata, Dana.

August 22–27, 1873. Mid-Atlantic, about 2° north of the Equator.

Calanus gracilis, Dana.

Eucalanus attenuatus, Dana.

Leuckartia ?

Undina vulgaris, Dana.

Scolecithrix danae (Lubbock).

Euchaeta prestandreae, Philippi.

Temora dubia (Lubbock).

Candace pachydactyla, Dana.

Corycaeus speciosus, Dana.

Copilia mirabilis, Dana.

Saphirinella stylifera (Lubbock).

August 25, 1873. Lat. $1^{\circ} 47'$ N., long. $24^{\circ} 26'$ W. 40 fathoms.

Undina vulgaris, Dana.

Euchaeta prestandreae, Philippi.