

*Alebion*, Kröyer.*Alebion*, Kröyer, Bidrag til Kundskab om Snyltekrebs, 1863.

Rostrum intermediate in form between that of *Caligus* and *Pandarus*, provided with large and strong palps. First pair of swimming feet two-branched, branches two-jointed; outer branches of the first, second, and third pairs armed with curiously-formed laminæ. Fourth pair rudimentary, two-jointed, one-branched. Carapace produced backwards into two long, narrow processes. Antennal palps, sucking-discs, and furcula wanting.

*Alebion carchariae*, Kröyer.

Kröyer, Bidrag til Kundskab om Snyltekrebs, p. 165, Taf. xii. fig. 1.

Found on *Zygæna malleus* at St. Vincent, Cape Verde.*Nogagus*, Leach.

This genus differs from *Pandarus* in that the cephalothoracic segments are destitute (or nearly so) of lateral alæ, and are not bilobate; the abdomen is well developed, and its segments distinct; there is no rounded telson, and the caudal stylets are laminar, terminal, and bear setæ of the true Cyclopoid type.

1. *Nogagus grandis*, Steenstr. and Ltk.*Nogagus grandis*, St. and Stk., Snyltekrebs og Lernæer, p. 45, Pl. x. fig. 19.

One specimen of this fine species. The memorandum of locality lost.

2. *Nogagus lunatus*, Steenstr. and Ltk.*Nogagus lunatus*, ibid. ibidem, p. 49, pl. ix. fig. 17.From back of *Carcharias glaucus*, Atlantic, March 2, 1876.3. *Nogagus latreillii*, Leach (*fide* Steenstr. and Ltk.).*Nogagus latreillii*, ibid. ibidem, p. 44, pl. ix. fig. 18.On *Carcharias brachyurus*, between Papua and Japan, and near the Kermadec Islands.4. *Nogagus curticaudis* (Dana).*Specilligus curticaudis*, Dana, Crust. U. S. Expl. Exped., p. 1375, pl. xcv. fig. 6.On *Carcharias brachyurus*, between Papua and Japan.5. *Nogagus validus*, Dana.*Nogagus validus*, idem. ibidem, p. 1363, pl. xciv. fig. 9.On *Carcharias brachyurus*, between Papua and Japan.