

## PLATE VI.

Figure 1.—*Leucilla connexiva*, n. sp.

|     |                                  | Diam. |
|-----|----------------------------------|-------|
| (a) | Portion of a horizontal section, | x 50  |
| (b) | A dermal triradiate spicule,     | x 100 |
| (c) | A gastric triradiate spicule,    | x 100 |
| (d) | A subgastric triradiate spicule, | x 100 |
| (e) | A subdermal triradiate spicule,  | x 100 |

,, 2.—*Leucilla uter*, n. sp.

|        |   |        |
|--------|---|--------|
| (a)    | Portion of a horizontal section ; <i>Am.</i> = Amphiblastulæ, | x 50   |
| (b)    | A dermal quadriradiate spicule,                               | x 100  |
| (c)    | A gastric quadriradiate spicule,                              | x 100  |
| (d, e) | Two subgastric triradiate spicules,                           | x 100  |
| (f)    | Two spermospores,   | x 1000 |

,, 3.—*Leuconia multiformis*, n. sp.

|     |                                  |       |
|-----|----------------------------------|-------|
| (a) | Portion of a horizontal section, | x 30  |
| (b) | Piece of the dermal surface,     | x 100 |
| (c) | A conjectural calcoblast,        | x 800 |