

6. *Discodoris muta*, Bergh.  
Antilles Sea.
7. *Discodoris modesta*, Bergh.  
Philippine Sea.
8. *Discodoris schmeltziana* (Garrett), Bergh.  
Pacific Ocean.
9. *Discodoris indecora*, Bergh.  
Mediterranean.
10. *Discodoris pardalis* (Alder and Hancock).  
Indian Ocean.
11. *Discodoris concinna* (Alder and Hancock).  
Indian Ocean.
12. *Discodoris fragilis* (Alder and Hancock).  
Indian Ocean.
13. *Discodoris vestita* ? (Abraham).  
Pacific Ocean, Straits of Magellan.
14. *Discodoris raripilosa* ? (Abraham).  
Habitat ?
15. *Discodoris stragulata* ? (Abraham).  
Habitat ?

*Discodoris morphaea*, Bergh (Pl. I. figs. 19–22; Pl. II. figs. 1–12; Pl. III. fig. 13).

*Discodoris morphaea*, Bergh, Malacolog. Untersuch., loc. cit., Heft xii., 1877, pp. 536–539,  
Taf. lx. figs. 18–22; Taf. lxi. figs. 1–5.

*Habitat*.—Philippine Sea (Samboangan).

A single individual of this species was dredged (along with a specimen of *Ceratosoma cornigerum* and one of *Platydoris eurychlamys*), February 1, 1875, on the reefs near Samboangan, at a depth of 10 fathoms.

The individual, which was very well preserved in alcohol, was 6·2 cm. long by 6 cm. broad and 1·8 cm. high; the breadth of the border of the mantle 2·3 cm., of the foot 1·8 cm.; the height of the rhinophores 4·5 mm.; the breadth of the (contracted) branchial fissure 4 mm., the height of the gill 8 mm.—The colour of the dorsal side was