

C O N T E N T S.

| | PAGE |
|---|--------|
| DISTRIBUTION of the NUDIBRANCHIATE GASTEROPODA in the Oceans, | 1 |
| LIST of NUDIBRANCHIATA dredged by H.M.S. Challenger, | 2 |
| I. NUDIBRANCHIATA KLADOHEPATICA, Bergh, | 2- 51 |
| Family PHYLLIROIDÆ, | 3- 8 |
| <i>Phylliroë</i> , Péron et Lesueur, | 4- 6 |
| <i>atlantica</i> , Bergh, | 5- 6 |
| <i>bucephala</i> , Péron et Lesueur, | 5 |
| <i>Acura</i> , H. and A. Adams, | 7- 8 |
| <i>pelagica</i> , Adams, | 7- 8 |
| <i>lanceolata</i> , Bergh, | 8 |
| Family ÆOLIDIADÆ, | 8- 30 |
| <i>Fiona</i> , Alder and Hancock, | 8- 10 |
| <i>marina</i> (Forskål), | 9- 10 |
| (<i>glandulæ ptyalinæ</i>), | 10-124 |
| <i>Glaucus</i> , Forster, | 10- 18 |
| <i>atlanticus</i> , Forster, | 16- 18 |
| <i>longicirrus</i> , Reinhardt, Bergh, | 18 |
| <i>Glaucilla briareus</i> (Reinhardt), Bergh, | 18 |
| <i>Janolus</i> , Bergh, | 18- 23 |
| <i>australis</i> , Bergh, | 19- 23 |
| <i>Cuthonella</i> , Bergh, | 23- 26 |
| <i>abyssicola</i> , Bergh, | 24- 26 |
| (<i>Splanchnotrophus</i> , Alder and Hancock), | 26 |
| <i>Rizzolia</i> , Trinchese, | 27- 30 |
| <i>australis</i> , Bergh, | 27- 30 |
| Family SCYLLEADÆ, | 31- 34 |
| <i>Scyllæa pelugica</i> , Linné, | 33- 34 |
| Family BORNELLIDÆ, | 34- 43 |
| <i>Bornella excepta</i> , Bergh, | 36- 43 |
| Family TRITONIADÆ, | 43- 51 |
| <i>Tritonia</i> , Cuvier, | 44- 47 |
| <i>challengeriana</i> , Bergh, | 45- 47 |
| <i>Marionia</i> , Vayssière, | 48- 51 |
| <i>occidentalis</i> , Bergh, | 49- 51 |