

Host	Species of <i>Myzostoma</i> .	Living.
<i>Actinometra meridionalis</i> , var. <i>quadrata</i> , P. H. C.	{ <i>M. areolatum</i> , n. sp.	Free.
,, <i>morsei</i> , P. H. C.	<i>M. nigrescens</i> , n. sp.	Do.
,, <i>mutabilis</i> , Lütken, MS.	{ <i>M. echinus</i> , n. sp.	Do.
,, <i>nigra</i> , Semp., MS.	{ <i>M. brevicirrum</i> , n. sp.	Do.
,, <i>nigra</i> , Semp., MS.	<i>M. brachiatum</i> , Graff.	Do.
,, <i>parvicirra</i> , Müll., sp.	{ <i>M. costatum</i> , F. S. L.	Do.
,, <i>parvicirra</i> , Müll., sp.	<i>M. pallidum</i> , Graff.	Do.
,, <i>parvicirra</i> , Müll., sp.	<i>M. triste</i> , Graff.	Do.
,, <i>parvicirra</i> , Müll., sp.	{ <i>M. elegans</i> , Graff.	Do.
,, <i>pulchella</i> , Pourt., sp.	{ <i>M. abundans</i> , n. sp.	Do.
,, <i>pulchella</i> , Pourt., sp.	{ <i>M. carinatum</i> , n. sp.	Do.
,, <i>pulchella</i> , Pourt., sp.	{ <i>M. inflator</i> , n. sp.	In independent sessile cysts.
,, <i>solaris</i> , Lam., sp.	<i>M. pallidum</i> , Graff.	Free.
,, <i>strata</i> , P. H. C.	<i>M. horologium</i> , n. sp.	Do.
<i>Comatula</i> , sp.?	<i>M. mæbianum</i> , n. sp.	Do.
,, <i>Comatula</i> , sp.?	{ <i>M. calycotyle</i> , n. sp.	Do.
,, <i>Comatula</i> , sp.?	{ <i>M. asymmetricum</i> , n. sp.	Sticking fast to and enlarging the pinnules.
,, <i>Comatula</i> , sp.?	<i>M. pentacrini</i> , n. sp.	In swellings on the arms.
,, <i>Comatula</i> , sp.?	{ <i>M. deformatior</i> , n. sp.	Inside the pinnules, which become swollen and ovoid.
,, <i>Comatula</i> , sp.?	{ <i>M. compressum</i> , n. sp.	Free.
,, <i>Comatula</i> , sp.?	{ <i>M. coronatum</i> , n. sp.	Do.
<i>Metacrinus costatus</i> , P. H. C.	<i>M. wyville-thomsoni</i> , n. sp.	Do.
,, <i>angulatus</i> , P. H. C.	<i>M. wyville-thomsoni</i> , n. sp.	Do.
<i>Hyocrinus</i> , sp.?	<i>Stelechopus hyocrini</i> , n. g., n. sp.	Do.