

In two out of the six series, the zoœcia are narrower than in the others, and apparently always uninhabited, and without any oral orifice.

Group B. FLUSTRINA.

Family IX. FLUSTRIDÆ.

Flustradæ, Bk., Brit. Mus. Cat., vol. i. p. 46; Gray (pars), Br. Rad., p. 145.

Flustridæ (pars) d'Orb., Rech. s. les Bryoz., Ann. d. Sci. Nat., 1851, p. 314; Hincks, Brit. Mar. Polyz. vol. i. p. 113; Smitt, Krit. Förteck. Skandinav. Hafsbryzoer, 1867, p. 357.

Escharidæ (pars), Johnst., Brit. Zooph., 1st ed. p. 248; 2nd ed. p. 263; d'Orb.

Polypiers à réseau (pars), Lamk.

Flustrées, Lamx., Exp. Méth.

Character.—Zoarium membranaceous, expanded and foliaceous, or ligulate; erect (sometimes decurrent on the support), uni- or bi-laminar. Zoœcia quincuncially arranged, more or less open and membranaceous in front, without a raised border; avicularia, when present, usually vicarious.

The Family here contains the following genera:—

§ A. Zoœcia contiguous.

§§ a. *Utrinque porosæ*, Linné.

1. *Flustra*, Linné.

- (1) *Flustra crassa*, n. sp. (Pl. XVI. fig. 6.)
- (2) *Flustra denticulata* Busk (Pl. XXXII. fig. 2).
- (3) *Flustra biseriata*, n. sp. (Pl. XVI. fig. 1).
- (4) *Flustra membraniporides*, n. sp. (Pl. XXXII. fig. 7).

§§ β. *Pagina altera tantum porosæ*, Linné.

2. *Carbasea*, Gray.

- (1) *Carbasea ovoidea*, Busk (Pl. XVI. fig. 3).
- (2) *Carbasea dissimilis*, Busk.
- (3) *Carbasea elegans*, Busk (Pl. XVI. fig. 5).
- (4) *Carbasea pedunculata*, n. sp. (Pl. XVI. fig. 4).
- (5) *Carbasea moseleyi*, n. sp. (Pl. XXXIII. fig. 4).
- (6) *Carbasea pisciformis*, Busk.
- (7) *Carbasea cribriformis*, Busk. (Pl. XXXIV. fig. 8).

§ B. Zoœcia disjunct.

3. *Diachoris*, Busk.

- (1) *Diachoris magellanica*, Busk.
var. *a distans* (Pl. XVI. fig. 2).