

PLATE XX.<sup>1</sup>

SMITTIA.—ADEONELLA.—PORELLA.—ESCHARA, &amp;c

|  | PAGE |
|--|------|
| Figure 1.— <i>Smittia tenuis</i> , . . . . .   | 150  |
| 1 <i>a</i> , magnified ; 1 <i>b</i> , single zoœcium, more highly magnified.   |      |
| „ 2.— <i>Adeonella regularis</i> (woodcut 53), . . . . .   | 186  |
| 2 <i>a</i> , magnified ; 2 <i>b</i> , single zoœcium, more highly magnified.   |      |
| „ 3.— <i>Porella laevis</i> , var. <i>subcompressa</i> , . . . . .   | 149  |
| 3 <i>a</i> , magnified ; 3 <i>b</i> , single zoœcium, more highly magnified.   |      |
| „ 4.— <i>Adeonella distoma</i> , var. <i>imperforata</i> , . . . . .   | 188  |
| 4 <i>a</i> , magnified ; 4 <i>b</i> , single zoœcium, more highly magnified.   |      |
| „ 5.— <i>Mucronella pyriformis</i> , . . . . .   | 155  |
| 5 <i>a</i> , 5 <i>b</i> , magnified.   |      |
| „ 6.— <i>Eschara elegantula</i> , . . . . .  | 141  |
| 6 <i>a</i> , 6 <i>b</i> , magnified.   |      |
| „ 7.— <i>Adeonella atlantica</i> (Pl. XXI. fig. 1) (woodcut 52), . . . . .   | 186  |
| 7 <i>a</i> , magnified ; 7 <i>b</i> , zoœcia, more highly magnified.   |      |
| „ 8.— <i>Flustramorpha marginata</i> , . . . . .   | 135  |
| 8 <i>a</i> , magnified ; 8 <i>b</i> , more highly magnified.   |      |
| „ 9.— <i>Mucronella contorta</i> , . . . . .   | 155  |
| 9 <i>a</i> , magnified ; 9 <i>b</i> , with avicularia ; 9 <i>c</i> , 9 <i>d</i> , orifice and avicularia, more highly magnified. |      |
| „ 10.— <i>Schizoporella tenuis</i> , . . . . .   | 165  |
| 10 <i>a</i> , magnified.   |      |

<sup>1</sup> The generic names given above have been substituted for that of *Eschara*, which, as seen in the text, I have now employed in a much more restricted sense.