

PLATE XXXIV.

PASYTHEA.—GEPHYROPHORA.—BRETTIA.—DIACHORIS, &c.

	PAGE
Figure 1.— <i>Pasythea eburnea</i> ,	5
1 <i>a</i> , magnified; 1 <i>b</i> , 1 <i>f</i> , erect zoecia; 1 <i>c</i> , adnate zoecium; 1 <i>d</i> , zoecia with radical tubes; 1 <i>e</i> , punctum, more highly magnified.	
„ 2.— <i>Gephyrophora polymorpha</i> ,	167
2 <i>a</i> , 2 <i>b</i> , magnified; 2 <i>c</i> , orifice.	
„ 3.— <i>Brettia australis</i> ,	7
„ 4.— <i>Diachoris costata</i> ,	60
4 <i>a</i> , articulated avicularium.	
„ 5.— <i>Phylactella</i> , sp. (?), fragment from Station 253, 3125 fathoms (not noticed in the text).	
„ 6.— <i>Brettia cornigera</i> ,	7
6 <i>a</i> , magnified; 6 <i>b</i> , lower internodes; 6 <i>c</i> , back view; 6 <i>d</i> , upper zoecium, more highly magnified.	
„ 7.— <i>Cellepora albirostris</i> (Pl. XXXV. fig. 3),	193
7 <i>a</i> , 7 <i>b</i> , magnified; 7 <i>c</i> , 7 <i>d</i> , avicularia; 7 <i>e</i> , operculum.	
„ 8.— <i>Carbasea cribriformis</i> , natural size,	58