

STATION 24, lat.  $18^{\circ} 38'$  N., long.  $65^{\circ} 5'$  W., off Culebra Island; 390 fathoms; Pteropod ooze.

<i>Pasythea eburnea</i> , B.		<i>Farciminaria atlantica</i> .
------------------------------	--	---------------------------------

CAPE VERDE ISLANDS, 100 to 120 fathoms.

<i>Retepora imperati</i> .		<i>Smittia jacobensis</i> .
<i>Porella laevis</i> , var. <i>subcompressa</i> .		

STATION 48, lat.  $43^{\circ} 4'$  N., long.  $64^{\circ} 5'$  W.; 51 fathoms; rock.

<i>Eschara elegantula</i> .		<i>Cellepora canaliculata</i> .
-----------------------------	--	---------------------------------

STATION 36, lat.  $32^{\circ} 7'$  N., long.  $65^{\circ} 4'$  W., off Bermuda; 30 fathoms; coral.

<i>Ætea anguina</i> , C, D, G.		<i>Bugula neritina</i> .
--------------------------------	--	--------------------------

CAPE VERDE ISLANDS, 10 fathoms.

<i>Diachoris hirtissima</i> .		<i>Cupularia owenii</i> .
-------------------------------	--	---------------------------

STATION 109, lat.  $0^{\circ} 55'$  N., long.  $29^{\circ} 22'$  W., off St. Paul's Rocks; shallow water.

<i>Catenaria diaphana</i> .		<i>Scrupocellaria macandrei</i> .
<i>Tubucellaria opuntioides</i> .		

#### B.—SOUTH ATLANTIC REGION.

STATION 325, lat.  $36^{\circ} 44'$  S., long.  $46^{\circ} 16'$  W.; 2650 fathoms; blue mud.

<i>Cellularia crateriformis</i> .		<i>Kinetoskias cyathus</i> , A.
<i>Farciminaria magna</i> , C.		

STATION 332, lat.  $37^{\circ} 29'$  S., long.  $27^{\circ} 31'$  W.; 2200 fathoms; Globigerina ooze.

<i>Bicellaria navicularis</i> .		<i>Bugula margaritifera</i> .
---------------------------------	--	-------------------------------

STATION 323, lat.  $35^{\circ} 39'$  S., long.  $50^{\circ} 47'$  W.; 1900 fathoms; blue mud.

<i>Cellularia crateriformis</i> .		<i>Farciminaria cribraria</i> .
<i>Bugula margaritifera</i> .		<i>Farciminaria magna</i> , var. <i>armata</i> .
<i>Salicornaria magnifica</i> , A, C.		