

STATION 315, lat.  $51^{\circ} 40'$  S., long.  $57^{\circ} 50'$  W.; 12 fathoms; sand and gravel.

<i>Menipea benemunita</i> , C, G.	<i>Microporella malusii</i> .
<i>Menipea aculeata</i> , G.	<i>Lepralia margaritifera</i> .
<i>Caberea minima</i> .	<i>Lepralia marsupium</i> .
<i>Carbasea ovoidea</i> , C, G.	<i>Chorizopora hyalina</i> , var. <i>bougainvillei</i> , C.
<i>Diachoris magellanica</i> , D.	<i>Smittia stigmatophora</i> .
<i>Diachoris costata</i> .	<i>Mucronella tricuspis</i> , C.
<i>Salicornaria malvinensis</i> , C, D, G.	<i>Cellepora eatonensis</i> , C, G.
<i>Cribrilina monoceros</i> , C, D, F, G.	

PORt WILLIAM, Falkland Islands; 5 to 10 fathoms.

<i>Carbasea ovoidea</i> , C, G.		<i>Salicornaria malvinensis</i> , C, D, G.
---------------------------------	--	--

### C.—SOUTH INDIAN OR KERGUELEN REGION.

STATION 156, lat.  $62^{\circ} 26'$  S., long.  $95^{\circ} 44'$  E.; 1975 fathoms; Diatom ooze.

*Bicellaria infundibulata*.

STATION 157, lat.  $53^{\circ} 55'$  S., long.  $108^{\circ} 35'$  E.; 1950 fathoms; Diatom ooze.

<i>Bugula bicornis</i> .		<i>Salicornaria magnifica</i> , A, B.
<i>Onchopora sinclairii</i> .		

STATION 153, lat.  $65^{\circ} 42'$  S., long.  $79^{\circ} 49'$  E.; 1675 fathoms; blue mud.

*Farciminaria magna*, B.

STATION 147, lat.  $46^{\circ} 16'$  S., long.  $48^{\circ} 27'$  E.; 1600 fathoms; Diatom ooze.

<i>Bicellaria infundibulata</i> .		<i>Bugula reticulata</i> , B, G.
<i>Foveolaria orbicularis</i> .		

STATION 148, lat.  $46^{\circ} 47'$  S., long.  $51^{\circ} 37'$  E., off Possession Island; 210 fathoms; hard ground, gravel, shells.

<i>Nellia oculata</i> , B, D, E.	<i>Smittia graciosa</i> .
<i>Caberea darwinii</i> , B.	<i>Mucronella ventricosa</i> , var. <i>multispinata</i> .
<i>Retepora gigantea</i> .	<i>Myriozoum marionense</i> .
<i>Retepora cavernosa</i> .	<i>Cellepora vagans</i> .
<i>Reteporella myriozoides</i> .	<i>Cellepora mamillata</i> , var. <i>atlantica</i> , B.
<i>Escharoides occlusa</i> , D, E.	<i>Schizoporella elegans</i> , B.