

TABLE V.—Showing the Association of Crinoid Genera at different Stations occupied by the "Porcupine," Challenger, and "Talisman," together with the number of Species obtained at each Station.

REMARKS.	Stations.	<i>Pentacrinus</i> .	<i>Metacrinus</i> .	<i>Rhizocrinus</i> .	<i>Bathyrinus</i> .	<i>Hyocrinus</i> .	<i>Antedon</i> .	<i>Actinometra</i> .	<i>Eudiocrinus</i> .	<i>Atelecrinus</i> .	<i>Promachocrinus</i> .	<i>Thaumatocrinus</i> .
Separate hauls at very different depths.	"Porcupine," 1870.	1	...	...	...	...	2	...	...	...	...	...
<i>Metacrinus murrayi</i> and <i>Pentacrinus naresianus</i> referred here conjecturally (see p. 128).	Challenger Stations. 210	1?	1?	...	...	...	2	...	...	...	...	...
... ..	175	1	...	...	...	...	3	...	...	...	...	...
... ..	235	1	...	...	...	...	...	...	1	...	...	...
... ..	170	1	2	...	...	...	6	...	...	...	...	...
Separate hauls at slightly different depths.	122	1	...	1	...	...	...	...	...	1	...	...
... ..	171	2	...	...	...	...	...	...	...	...	...	...
... ..	214	2	4	...	...	...	6	...	...	...	1	...
... ..	192	...	4	...	...	...	12	1	...	...	...	...
<i>Hyocrinus</i> , stem only ( <i>fide</i> C.W.T.).	106	...	...	...	1	1?	...	...	...	...	...	...
... ..	147	...	...	...	1	1	3	...	...	...	1	...
... ..	158	...	...	...	...	...	...	...	...	...	1	1
No information about Comatulæ.	{ "Talisman," 1882.	...	...	1	...	...	...	...	...	...	...	...
	1883. { A.	1	...	...	...	...	...	...	...	...	...	...
	{ B.	...	...	1	1	...	...	...	...	...	...	...

Table VI. shows the association of *Pentacrinus*, *Holopus*, *Rhizocrinus*, and the three Comatulid genera in the Caribbean Sea, together with the number of times that two or more species of *Pentacrinus* were found at the same Station. Comatulæ were obtained some eighty times; but as I have not yet been able to determine them critically, I have found it impossible to give the number of species occurring at each Station, as has been done for the Challenger dredgings.