

TABLE VI.—Showing the Association of Crinoid Genera in the Caribbean Sea, together with the number of times that two or more Species of *Pentacrinus* were found at the same Station.

Number of Stations.	<i>Pentacrinus</i> .	<i>Rhizocrinus</i> .	<i>Holopus</i> .	<i>Antedon</i> .	<i>Actinometra</i> .	<i>Atelecrinus</i> .
13	1 species					
12		×				
1	1 "	×				
1	1 "	×		×	×	
3	1 "			×		
3	1 "			×	×	
1	1 "				×	
4	2 "					
1	2 "	×		×		
1	2 "	×			×	
1	2 "			×		
2	2 "			×	×	
1	4 "		×	×	×	
2			×			
1		×		×		
4		×			×	
3		×		×	×	
1		×				×
1		×		×		×
Stalked Crinoids at 56 Stations.	<i>Pentacrinus</i> at 32 Stations.	<i>Rhizocrinus</i> at 26 Stations.	<i>Holopus</i> at 3 Stations (two isolated).	Comatulæ at 82 Stations altogether.		

The above Table embraces—(1) all the dredgings of the U. S. Coast Survey Steamers in the Caribbean Sea and Gulf Stream; (2) those of Captain Cole of the Telegraph Steamer "Investigator;" (3) those carried on at Barbadoes by the collectors for Sir Rawson Rawson; (4) those of a few isolated collectors who have sent specimens to the Museum of Comparative Zoology. The whole series will be found recorded in the Station List at the end of the Report. No attempt has been made to classify any observations anterior to those of Pourtalès in 1867.