

STATION 158. March 7, 1874; lat. $50^{\circ} 1' S.$, long. $123^{\circ} 4' E.$; 1800 fathoms; Globigerina ooze; bottom temperature, $33^{\circ} \cdot 5 F.$

Thaumatocrinus renovatus.

Promachocrinus abyssorum. Also at Station 147.

STATION 170A. July 14, 1874; near the Kermadec Islands; lat. $29^{\circ} 45' S.$, long. $178^{\circ} 11' W.$; 630 fathoms; volcanic mud; bottom temperature, $39^{\circ} \cdot 5 F.$

Pentacrinus naresianus. Also at Stations 171, 175, 210 (?), 214.

Metacrinus nodosus

Metacrinus wyvillii. Also at Station 214.

Antedon (six species).

STATION 171. July 15, 1874; off Raoul Island; lat. $28^{\circ} 33' S.$, long. $177^{\circ} 50' W.$; 600 fathoms; hard ground; bottom temperature, $39^{\circ} \cdot 5 F.$

Pentacrinus alternicirrus. Also at Stations 210 (?), 214.

Pentacrinus naresianus. Also at Stations 170A, 175, 210 (?), 214.

STATION 175. August 12, 1874; near Kandavu, Fiji; lat. $19^{\circ} 2' S.$, long. $177^{\circ} 10' E.$; 1350 fathoms; Globigerina ooze; bottom temperature, $36^{\circ} F.$

Pentacrinus naresianus. Also at Stations 170A, 171, 210 (?), 214.

Antedon (three species).

STATION 192. September 26, 1874; near the Ki Islands; lat. $5^{\circ} 49' S.$, long. $132^{\circ} 14' E.$; 140 fathoms; blue mud.

Metacrinus angulatus.

Metacrinus cingulatus.

Metacrinus murrayi (?).

Metacrinus nobilis.

Metacrinus tuberosus.

Antedon (twelve species).

Actinometra (one species).

STATION 209. January 22, 1875; lat. $10^{\circ} 14' N.$, long. $123^{\circ} 54' E.$; 95 fathoms; blue mud; bottom temperature, $71^{\circ} F.$

Metacrinus interruptus.