

PLATE XXIII.

- Fig. 1. Ventral aspect of the anterior region of *Eulepis challengerica*, . Enlarged.
- Fig. 2. Ventral view of the anterior region of *Psammolyce occidentalis*, Enlarged.
- Fig. 3. Dorsal aspect of the head and anterior region of the same.
The palpi have no connection with the first feet, which
lie above them.
- Fig. 4. Anterior region of *Thalenessa fimbriata*, viewed from the
dorsum, Enlarged.
- Fig. 5. Anterior dorsal region of *Thalenessa digitata*, Enlarged.
- Fig. 6. Foot of the same, × about 20.
- Fig. 7. Scale of the foregoing, × about 20.
- Fig. 8. Foot of *Eupompe australiensis*. From the front. The articula-
tion of the dorsal cirrus is almost hidden, × about 20.
- Fig. 9. Scale of *Leanira hystericis*, × about 20.
- Fig. 10. Scale of *Leanira lævis*, × about 20.
- Fig. 11. Foot from the anterior third of *Leanira lævis*, × about 25.
- Fig. 12. Papilla from scale of *Thalenessa oculata*, × 210.
- Fig. 13. Foot (mounted in Farrant's solution) from the anterior third of
Leanira magellanica, × about 20.