

PLATE XXXII.

- | | | | | | | |
|----------|--|---|---|---|---|-----------|
| Fig. 1. | Anterior region of <i>Eunoa iphionoides</i> , | . | . | . | . | Enlarged. |
| Fig. 2. | Head of <i>Eupolynoë mollis</i> , | . | . | . | . | Enlarged. |
| Fig. 3. | Head and anterior region of <i>Harmothoë haliaëti</i> , | . | . | . | . | Enlarged. |
| Fig. 4. | Portion of the surface of a scale of the same, | . | . | . | . | × 55. |
| Fig. 5. | Edge of a posterior scale of the same, | . | . | . | . | × 55. |
| Fig. 6. | Head and anterior region of <i>Polynoë ascidioides</i> , | . | . | . | . | Enlarged. |
| Fig. 7. | Posterior border of a scale of <i>Dasylepis equitis</i> , | . | . | . | . | × 55. |
| Fig. 8. | Thirtieth foot of <i>Phyllodoce (Anaitis) sanctæ vincentis</i> , | . | . | . | . | × 40. |
| Fig. 9. | Thirtieth foot of <i>Phyllodoce duplex</i> , | . | . | . | . | × 24. |
| Fig. 10. | Thirtieth foot of <i>Eulalia capensis</i> , | . | . | . | . | × 24. |
| Fig. 11. | Foot of <i>Greeffia oahuensis</i> , | . | . | . | . | × 18. |
| Fig. 12. | Middle foot of <i>Alciopa antarctica</i> , from Honolulu, | . | . | . | . | × 18. |
| Fig. 13. | Foot of <i>Alciopa antarctica</i> , variety, | . | . | . | . | × 24. |
| Fig. 14. | Anterior foot of <i>Hesione pacifica</i> , | . | . | . | . | × 25. |
| Fig. 15. | Foot of <i>Hesione</i> (fragment), from St. Thomas, West Indies, | . | . | . | . | × 25. |
| Fig. 16. | Foot of <i>Hesione (Fallacia) pantherina</i> , | . | . | . | . | × 25. |