

PLATE XXXIV.

Fig. 1. Anterior region of <i>Syllis robertianaæ</i> ,	.	.	.	.	x about 25.
Fig. 2. Anterior foot of the foregoing,	.	.	.	.	x 55.
Fig. 3. Head and anterior region of <i>Nereis kobiensis</i> ,	.	.	.	.	Enlarged.
Fig. 4. Tenth foot of the foregoing,	.	.	.	.	x 25.
Fig. 5. Thirty-seventh foot of the same,	.	.	.	.	x 25.
Fig. 6. Maxilla of the same species,	.	.	.	.	x 25.
Fig. 7. Head and anterior region of <i>Nereis tongatabuensis</i> ,	.	.	.	.	Enlarged.
Fig. 8. Tenth foot of the same,	.	.	.	.	x 25.
Fig. 9. Thirty-seventh foot of the foregoing,	.	.	.	.	x 25.
Fig. 10. Head and anterior region of <i>Nereis arafurensis</i> . The palpi have been bent downward and inward,	.	.	.	.	Enlarged.
Fig. 11. Tenth foot of the same,	.	.	.	.	x 25.
Fig. 13. Thirty-seventh foot of the foregoing,	.	.	.	.	x 25.
Fig. 14. Head and anterior region of <i>Nereis melanocephala</i> ,	.	.	.	.	Enlarged.
Fig. 15. Tenth foot of the same,	.	.	.	.	x 25.
Fig. 16. Thirty-seventh foot of the same,	.	.	.	.	x 25.
Fig. 17. Sixty-seventh foot of the foregoing,	.	.	.	.	x 40.