

PLATE XXXV.

Fig. 1. Head and anterior region of <i>Nereis atlantica</i> ,	:	Enlarged.
Fig. 2. Tenth foot of the foregoing,	:	× 40.
Fig. 3. Fifty-seventh foot of the same,	:	× 40.
Fig. 4. Head and anterior region of <i>Nereis longisetis</i> ,	:	Enlarged.
Fig. 5. Head and anterior region of <i>Nereis eatoni</i> ,	:	Enlarged.
Fig. 6. Proboscis of the same, with maxillæ and paragnathi. From a small specimen,	:	Enlarged.
Fig. 7. Head and anterior region of <i>Nereis antillensis</i> ,	:	Enlarged.
Fig. 8. Tenth foot of the same,	:	× 40.
Fig. 9. Twenty-sixth foot of the foregoing,	:	× 40.
Fig. 10. Head, proboscis, and anterior region of <i>Nereis kerguelensis</i> ,	:	Enlarged.
Fig. 11. Tenth foot of the same,	:	× 25.
Fig. 12. Thirty-seventh foot of the foregoing,	:	× 25.
Fig. 13. Head and anterior region of <i>Nereis patagonica</i> . The terminal bosses of the palpi are doubled under,	:	Enlarged.
Fig. 14. Tenth foot of a small example of the foregoing,	:	× 40.
Fig. 15. Thirty-seventh foot of a large example of the same,	:	× 25.