

PLATE XXXVI.

Fig. 1. Head and anterior end of <i>Nereis brasiliensis</i> ,	Enlarged.
Fig. 2. Tenth foot of the same,	× 25.
Fig. 3. Thirty-seventh foot of the foregoing,	× 25.
Fig. 4. Anterior foot of <i>Staurocephalus atlanticus</i> , indifferently preserved,	× 80.
Fig. 5. Head and anterior region of the same species,	Enlarged.
Fig. 6. Foot of <i>Staurocephalus australiensis</i> ,	× 25.
Fig. 7. Anterior region (dorsal view) of <i>Lumbriconereis pettigrewi</i> ,	Enlarged.
Fig. 8. Tenth foot of the same,	× 40.
Fig. 9. Thirtieth foot of the foregoing,	× 40.
Fig. 10. Dorsal aspect of the anterior region of <i>Lumbriconereis bifurcata</i> ,	Enlarged.
Fig. 11. Ventral view of the anterior region of the same species,	Enlarged.
Fig. 12. Posterior foot of the foregoing,	× 40.
Fig. 13. Dorsal view of the anterior region of <i>Lumbriconereis japonica</i> ,	Enlarged.
Fig. 14. Tenth foot of the same,	× 25.
Fig. 15. Posterior foot of the foregoing, seen partly from below,	× 25.
Fig. 16. Dorsal aspect of the anterior region of <i>Lumbriconereis kerguelensis</i> ,	Enlarged.
Fig. 17. Fiftieth foot of the foregoing,	× 80.
Fig. 18. Dorsal view of the anterior region of <i>Lumbriconereis neo-zealandiae</i> . The longitudinal grooves on the dorsum have not been shown,	Enlarged.
Fig. 19. Thirty-third foot of the same, seen obliquely from below,	× 80.
Fig. 20. Dorsal view of the anterior region of <i>Lumbriconereis abyssorum</i> ,	Enlarged.
Fig. 21. Thirtieth foot of the same,	× 80.