

PLATE XLII.

Fig. 1. Anterior dorsal region of <i>Nothria ehlersi</i> ,	Enlarged.
Fig. 2. Tenth foot of the same,	× 25.
Fig. 3. Thirtieth foot of the foregoing,	× 25.
Fig. 4. Foot of <i>Eone trifida</i> ,	× 80.
Fig. 5. Thirtieth foot of <i>Glycera tessellata</i> ,	× 40.
Fig. 6. Twentieth foot of <i>Glycera amboinensis</i> ,	× 80.
Fig. 7. Anterior region of the same, with the proboscis extruded,	× about 12.
Fig. 8. Thirtieth foot of <i>Glycera sagittariae</i> ,	× 40.
Fig. 9. Anterior region of <i>Glycera lamelliformis</i> , with the proboscis ex- truded,	Slightly enlarged.
Fig. 10. Thirtieth foot of <i>Glycera lamelliformis</i> ,	× 15.
Fig. 11. Dorsal view of the anterior region of <i>Hemipodus magel- lanicus</i> ,	Somewhat enlarged.
Fig. 12. Ventral view of the same region in another example,	Similarly enlarged.
Fig. 13. Tenth foot of the same,	× 25.
Fig. 14. Thirtieth foot of the foregoing,	× 25.
Fig. 15. One hundred and tenth foot of <i>Hemipodus magellanicus</i> ,	× 25.