

PLATE VIa.

Fig. 1.	Tip of a ventral bristle of <i>Aphrodita intermedia</i> ,	× 210.
Fig. 2.	Dorsal spine of <i>Aphrodita echidna</i> ,	× 18.
Fig. 3.	Tip of one of the lower ventral bristles of the same species,	× 210.
Fig. 4.	Tip of a light bronze-coloured dorsal spine of <i>Aphrodita australis</i> ,	× 210.
Fig. 5.	Tip of one of the inner dorsal bristles (which curve inward amongst the felt),	× 350.
Fig. 6.	Tip of one of the smaller inferior ventral bristles of the same species,	× 102.
Fig. 7.	Extremity of one of the dart-shaped anterior ventral bristles, the tip being broken,	× 350.
Fig. 8.	Spinose dorsal bristle of <i>Palmyra aurifera</i> ,	× 50.
Fig. 9.	Ventral bristle of the same,	× 210.
Fig. 10.	Dorsal bristle of <i>Lagisca antarctica</i> in Farrant's solution,	× 90.
Fig. 11.	Ventral bristle of the same,	× 90.
Fig. 12.	Dorsal bristle of <i>Evarne kerguelensis</i> ,	× 210.
Fig. 13.	Superior ventral bristle of the same, slightly turned round, and showing a trace of a secondary process,	× 210.
Fig. 14.	One of the series having shorter tips from the middle of the ventral tuft in the foregoing species,	× 210.