

PLATE VIII A.

Fig. 1. Tip of one of the stouter dorsal bristles of <i>Eunoa iphionoides</i> ,	× 210.
Fig. 2. Ventral bristle of the same species,	× 210.
Fig. 3. Tip of one of the attenuate silky hairs of the dorsal tuft of <i>Iphione spinosa</i> ,	× 700.
Fig. 4. Inferior portion of the same bristle,	× 350.
Fig. 5. Tip of a ventral bristle of the same species in profile,	× 90.
Fig. 6. A small bristle from the same region of the foot, seen antero-posteriorly,	× 90.
Fig. 7. One of the fine pale dorsal bristles of <i>Iphione ovata</i> ,	× 700.
Fig. 8. Tip of one of the second ventral series in the same species,	× 350.
Fig. 9. Tip of a dorsal bristle of <i>Eunoa opalina</i> ,	× 90.
Fig. 10. Abnormal tip (from fracture ?) of one of the same bristles,	× 90.
Fig. 11. Pale yellow ventral bristle of the foregoing,	× 90.
Fig. 12. Dorsal bristle of <i>Lænilla fusca</i> ,	× 90.
Fig. 13. One of the average examples, from the ventral division of the foot,	× 90.
Fig. 14. Tip of one of the longer superior ventral bristles of <i>Polynoë platycirrata</i> ,	× 210.
Fig. 15. Antero-posterior view of a smaller example from the same region,	× 210.