

PLATE IXA.

Fig. 1. Dorsal bristles of <i>Polynoë platycirrata</i> ,	× 350.
Fig. 2. Dorsal bristle of <i>Lepidonotus gymnonotus</i> ,	× 90.
Fig. 3. Tip of average ventral bristle of the same,	× 210.
Fig. 4. Dorsal bristle of <i>Polynoë capensis</i> ,	× 350.
Fig. 5. Ventral bristle of the same,	× 350.
Fig. 6. Dorsal bristle of <i>Polynoë pustulata</i> ,	× 350.
Fig. 7. Tip of a ventral bristle of the same form,	× 350.
Fig. 8. Attenuate dorsal bristle of <i>Polynoë (Langerhansia) euplectellæ</i> ,	× 350.
Fig. 9. Tip of average ventral bristle of the foregoing,	× 210.
Fig. 10. Tip of a dorsal bristle of <i>Evarne tenuisetis</i> ,	× 350.
Fig. 11. Extremity of one of the superior ventral bristles of the same species,	× 350.
Fig. 12. Tip of an inferior ventral bristle of the same,	× 350.
Fig. 13. Tip of dorsal bristle of <i>Lagisca magellanica</i> , var. <i>murrayi</i> ,	× 90.
Fig. 14. One of the short ventral bristles (from the middle of the series) of the same,	× 90.