

PLATE XIA.

- Fig. 1. One of the shorter and peculiarly curved dorsal bristles of *Eunoa capensis*, × 350.
- Fig. 2. Tip of another, with a longer smooth extremity, × 350.
- Fig. 3. Ventral bristle of the same, × 350.
- Fig. 4. One of the longer and more slender dorsal bristles of *Lagisca crosetensis*, × 90.
- Fig. 5. Lateral view of the tip of one of the shorter dorsal bristles of the same, × 350.
- Fig. 6. Average example from the ventral division of the foot in the foregoing, × 210.
- Fig. 7. Spine-like bristle from the ventral division of the foot of *Polynoella levisetosa*, × 90.
- Fig. 8. Dorsal bristle of *Polynoë attenuata*, × 350.
- Fig. 9. Tip of ventral bristle, from the same, × 350.
- Fig. 10. Tip of dorsal bristle of *Lepidonotus cristatus*, × 350.
- Fig. 11. Tip of a ventral bristle of the foregoing, × 210.
- Fig. 12. Dorsal bristle of *Lagisca yokohamiensis* (average example), × 350.
- Fig. 13. Ventral bristle of the same, × 210.
- Fig. 14. An average form amongst the longer dorsal bristles of *Eunoa abyssorum*, × 90.
- Fig. 15. Tip of one of the longer (superior) ventral bristles of the preceding, × 90.
- Fig. 16. Tip of one of the shorter and smaller inferior bristles, × 90.